

# Examination of the generic concept and species boundaries of the genus *Erioscyphella* (Lachnaceae, Helotiales, Ascomycota) with the proposal of new species and new combinations based on the Japanese materials

Yukito Tochihara<sup>1,2</sup>, Tsuyoshi Hosoya<sup>2</sup>

**1** Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan **2** Department of Botany, National Museum of Nature and Science, 4-1-1 Amakubo, Tsukuba, Ibaraki 305-0005, Japan

Corresponding author: Yukito Tochihara ([tochi@kahaku.go.jp](mailto:tochi@kahaku.go.jp))

---

Academic editor: Cecile Gueidan | Received 17 August 2021 | Accepted 10 January 2022 | Published 8 February 2022

---

**Citation:** Tochihara Y, Hosoya T (2022) Examination of the generic concept and species boundaries of the genus *Erioscyphella* (Lachnaceae, Helotiales, Ascomycota) with the proposal of new species and new combinations based on the Japanese materials. MycoKeys 87: 1–52. <https://doi.org/10.3897/mycokeys.87.73082>

---

## Abstract

The genus *Erioscyphella* Kirschst., which was morphologically confused with *Lachnum*, was herein examined. Based on molecular phylogenetic analyses using a combined dataset of ITS, LSU, mtSSU, and RPB2 and morphological examinations, *Erioscyphella* was distinguished from *Lachnum* and redefined by longer ascospores and the presence of apical amorphous materials and/or resinous materials equipped on hairs. Species boundaries recognized by morphology/ecology and phylogenetic analyses were cross-checked using species delimitation analyses based on DNA barcode sequences downloaded from UNITE, resulting in that species' taxonomic problems being uncovered. Six new species (*E. boninensis*, *E. insulae*, *E. otanii*, *E. papillaris*, *E. paralushanensis*, and *E. sasibrevispora*) and two new combinations (*E. hainanensis* and *E. sinensis*) were proposed.

## Keywords

ITS, morphology, phylogeny, species delimitation, species hypothesis, taxonomy, UNITE

## Introduction

The genus *Erioscyphella* Kirschst belongs to the family Lachnaceae Raitv. (Helotiales, Ascomycota) and includes 11 species: *E. abnormis* (Mont.) Baral, Šandová & B. Perić [lectotype of *Erioscyphella* (Haines and Dumont 1984); as '*E. longispora* (P. Karst.) Kirschst.' in the original description (Kirschstein 1938)], *E. alba* Ekanayaka & K.D. Hyde, *E. aseptata* Ekanayaka & K.D. Hyde, *E. bambusina* (Bres.) Kirschst., *E. brasiliensis* (Mont.) Baral, Šandová & B. Perić, *E. curvispora* B. Perić & Baral, *E. euterpes* (S.A. Cantrell & J.H. Haines) Guatim., R.W. Barreto & Crous, *E. fusiformis* (Ekanayaka & K.D. Hyde) Ekanayaka & K.D. Hyde, *E. lunata* (W.Y. Zhuang & Spooner) B. Perić & Baral, *E. lushanensis* (W.Y. Zhuang & Zheng Wang) Guatim., R.W. Barreto & Crous, and *E. sclerotii* (A.L. Sm.) Baral, Šandová & B. Perić. (Index Fungorum 2021).

*Erioscyphella* has been suggested as a monophyletic group by molecular phylogenetic analyses by Cantrell and Hanlin (1997), Hosoya et al. (2010), Perić and Baral (2014), and Guatimosim et al. (2016). However, the morphological delimitation of the genus is currently ill-defined. In the original description (Kirschstein 1938), *Erioscyphella* was misleadingly defined based on features that are not taxonomically informative, such as filiform, colored, and pigmented ascospores and lanceolate paraphyses (Korf 1978; Perić and Baral 2014). After that, in the genus *Lachnum* Retz. [type genus of Lachnaceae], species of so-called 'long-spored *Lachnum*', which were characterized by longer ascospores and the occurrence in tropical areas, were suggested as members of *Erioscyphella* (Haines and Dumont 1984) and have been transferred into *Erioscyphella* based on molecular phylogenetic analyses by Perić and Baral (2014) and Guatimosim et al. (2016). However, in fact, as morphology of *Erioscyphella*, including 'long-spored *Lachnum*', is consecutive with that of the genus *Lachnum* especially regarding the ascospore length and shape of ectal excipular cells (Haines and Dumont 1984), the morphological delimitation of *Erioscyphella* has not been sufficiently discussed. Since much more potential species are thought to be included in *Erioscyphella*, its morphological concept must be discussed and updated based on a wider size of taxon sampling.

In the present study, the authors aimed to: a) clarify the generic boundaries of *Erioscyphella* using molecular and morphological/ecological data, and b) propose new species or new combinations based on more objectively defined species boundaries. To reach our first goal, we used specimens from the herbarium of the National Museum of Nature and Science (TNS) (Tsukuba, Japan) as most of them were accompanied by culture and/or DNA extracts. In TNS, only three identified species of *Erioscyphella* were recognized (*E. abnormis*, *E. brasiliensis*, and *E. sclerotii*); however, we presumed that many unidentified species of *Erioscyphella* were housed therein. To reach our second goal, for species recognition, we tested DNA barcoding using the internal transcribed spacer region of nuclear ribosomal DNA (ITS), widely accepted as fungal DNA barcode (Begerow et al. 2010; Schoch et al. 2012; Hosoya 2021). ITS-based species boundaries were explored based on multiple methods, and the results were compared to species boundaries based on morphology, ecology, and phylogenetic relationships.

## Materials and methods

### Taxon sampling

In TNS, specimens labeled as *Erioscyphella* were initially searched, and closely related specimens to *Erioscyphella* were searched based on the sequence similarities of ITS. Selected specimens were tentatively identified based on morphology following Dennis (1954), Haines (1980), Haines and Dumont (1984), Spooner (1987), and Perić and Baral (2014).

### Morphological observation, DNA extraction, and sequencing

Micromorphology was examined using cotton blue (CB) dissolved in lactic acid (LA) (CB/LA; 0.5 g CB and 99.5 mL LA) as a mounting fluid. To check the ascal apex iodine reaction, Melzer's reagent (MLZ; 0.5 g I<sub>2</sub>, 1.5 g KI, 20 g chloral hydrate, and 20 g water) was initially used without KOH pretreatment, and Lugol's iodine (IKI; 1 g I<sub>2</sub> and 1 g KI, and 100 mL H<sub>2</sub>O) and MLZ with 3% KOH pretreatment were used when necessary. World Geodetic System 84 was used for the geographic coordinates. URLs herein shown were accessed on April 15, 2021, except for GBIF website accessed on Feb 10, 2020.

DNA was extracted from cultivated isolates in 2% malt extract broth (MEB) using the modified cetyltrimethylammonium bromide (CTAB) method (Hosaka and Castellano 2008; Tochiara and Hosoya 2019). When isolates are not available, DNA was extracted directly from a crushed apothecium using DNA extraction buffer following Tochiara and Hosoya (2019). The isolates from which DNA extracted were deposited in the NITE National Biological Resource Center (NBRC) (Kisarazu, Japan), except for isolates with restriction on transition by Japanese laws and those unavailable because of contracts with private companies.

Polymerase chain reaction (PCR) was used to amplify the following regions: ITS (= ITS1-5.8S-ITS2), the partial large subunit nuclear ribosomal RNA gene (LSU), the partial mitochondrial small subunit (mtSSU), and section '6–7' of the second largest subunit of the nuclear RNA polymerase II gene (RPB2). Primer pairs for PCR reactions of ITS, LSU and mtSSU were ITS1F (5'-CTTGGTCATTTAGAGGAAG-TAA-3') (Gardes and Bruns 1993) or ITS1 (5'-TCCGTAGGTGAACCTCGGG-3') (White et al. 1990) and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (White et al. 1990), LR0R (5'-ACCCGCTGAACTTAAGC-3') and LR5 (5'-TCCTGAGG-GAACTTCG-3') (Vilgalys and Hester 1990), and mrSSU1 (5'-AGCAGTGAG-GAATATTGGTC-3') and mrSSU3R (5'-ATGTGGCACGTCTATAGCCC-3') (Zoller et al. 1999) respectively. The PCR program consisted of an initial denaturation at 95 °C for 3 min, followed by 30 cycles of 94 °C for 35 s, 51 °C for 30 s, and 72 °C for 1 min, and a final extension at 72 °C for 10 min. When appropriate PCR products were not obtained, a modified PCR program was applied first, and then alternative primer pairs were tested. For RPB2, an alternative forward primer fRPB2-5F (5'-GAY-

GAYMGWGATCAYTTYGG–3') (Liu et al. 1999) or RPB2-P6Fa (5'–TGGGGGRYTK GTBTGYCCKGCHGA–3') (Hansen et al. 2005) and a reverse primer bRPB2-7.1R2 (5'–CCCATNGCYTGYTTVCCCATDGC–3') (modified from bRPB2-7.1R) (Matheny 2005; Matheny et al. 2007; Gelardi et al. 2015) were used.

Sequencing was conducted on an ABI PRISM 3500xL Genetic Analyzer (Applied Biosystems; Thermo Fisher Scientific, Waltham, MA, USA) with a BigDye Terminator 3.1 Cycle Sequencing Kit (Applied Biosystems). The obtained sequences were assembled using ATGC 7 (Genetyx, Tokyo, Japan). Assembled sequences were deposited in the International Nucleotide Sequence Database Collaboration (INSDC) via the DNA Data Bank of Japan (DDBJ), and acquired INSDC accession numbers. Assembled ITS sequences were also deposited in the UNITE database (<https://unite.ut.ee/>) via the PlutoF workbench (<https://plutof.ut.ee/>) (Abarenkov et al. 2010) and acquired UNITE accession numbers.

## Phylogenetic analyses

The specimens obtained from TNS were included in the phylogenetic analyses as candidate members of *Erioscyphella* ('‡' in Table 1). From other genera of the family Lachnaceae, four species of *Lachnum*, two species of *Albotricha*, *Brunnipila*, *Capitotricha*, *Dasyscyphella*, *Incrucipulum*, and *Lachnellula*, and one species of *Neodasyscypha* and *Proliferodiscus* were used ('†' in Table 1). Among the eight genera, seven of them (except *Proliferodiscus*) included type species. Three species of Helotiales were selected as outgroups following Tochihara and Hosoya (2019) (Table 1).

A concatenated dataset of ITS, LSU, mtSSU, and RPB2 was used in the phylogenetic analyses. Each region was aligned separately using MAFFT 7 (Kato and Standley 2013). The Q-INS-i option was used for ITS, LSU, and mtSSU to accommodate the secondary structures of RNA, and the G-INS-1 option was used for RPB2 to assume global alignment using the entire region. The aligned sequences were edited manually using BioEdit 7.0.5.2 (Hall 1999).

Phylogenetic conflicts among gene partitions were checked before the phylogenetic analyses using the concatenated matrix. Maximum likelihood (ML) trees with 1,000 bootstrap replications (Felsenstein 1985) using the ITS, LSU, mtSSU, and RPB2 datasets separately were constructed using MEGA X (Kumar et al. 2018) with the GTR+G model; branches with bootstrap values > 70% were compared among trees. For mtSSU and RPB2, specimens containing missing data were excluded from the analyses.

The concatenated dataset was analyzed using ML, maximum parsimony (MP), and Bayesian inference (BI). For the ML and BI analyses, substitution models were estimated for each partition (ITS, LSU, mtSSU, and each codon position of RPB2) based on Akaike's information criterion (AIC) (Akaike 1974) using Modeltest-NG 0.1.6 (Darriba et al. 2019).

ML tree search (Felsenstein 1984) and bootstrapping (Felsenstein 1985; Lemoine et al. 2018) was performed using RAxML-NG 0.9.0 (Kozlov et al. 2019) with 1,000 bootstrap replications under the substitution model SYM+I+G4 for ITS, TIM1+I+G4

Table 1. Taxa analyzed in the phylogenetic analyses.

| Specimen no.<br>(TNS-F-) | Taxon <sup>1</sup>                                                      | Collection site                                        | Collected<br>Date | Host plants and parts                    | Strain no.<br>(NBRC <sup>2</sup> ) | UNITE/GenBank accession no. <sup>3</sup> |          |          |          |
|--------------------------|-------------------------------------------------------------------------|--------------------------------------------------------|-------------------|------------------------------------------|------------------------------------|------------------------------------------|----------|----------|----------|
|                          |                                                                         |                                                        |                   |                                          |                                    | ITS                                      | LSU      | mtSSU    | RPB2     |
| †16740                   | <i>Albotricha acutipila</i> (P. Karst.) Raitv.                          | JAPAN, Nagano, Ueda, Sugadaira Montane Research Center | 2006-06-17        | culm of unidentified bamboo              | 104380                             | AB481234                                 | LC438571 | LC431751 | AB481354 |
| †16497                   | <i>Albotricha albotestacea</i> (Desm.) Raitv.                           | JAPAN, Nagano, Ueda, Sugadaira Montane Research Center | 2005-05-18        | culm of <i>Miscanthus sinensis</i>       | 101346                             | AB481235                                 | LC424943 | LC431747 | AB481340 |
| †16635                   | <i>Brunnipila fuscescens</i> (Pers.) Baral                              | JAPAN, Gunma, Higashi-Agatsuma                         | 2006-04-27        | leaf of unidentified tree                | 104365                             | AB481255                                 | LC424945 | LC431750 | AB481348 |
| †16690                   | <i>Brunnipila pseudocannabina</i> (Raitv.) Tochihara, Sasagawa & Hosoya | JAPAN, Akita, Kosaka                                   | 2006-05-26        | stem of unidentified herb                | 104374                             | AB481272                                 | LC533520 | LC533522 | LC533521 |
| †65670                   | <i>Capitotricha bicolor</i> (Bull.) Baral                               | SWITZERLAND, Filisur                                   | 2016-06-06        | twig of <i>Prunus spinosa</i>            | (FC-6101)                          | LC424834                                 | LC424942 | LC533244 | LC425011 |
| †65752                   | <i>Capitotricha rubi</i> (Bres.) Baral                                  | SWITZERLAND, Saicourt                                  | 2016-06-04        | twig of <i>Rubus idaeus</i>              | (FC-6075)                          | LC438560                                 | LC438573 | LC533243 | LC440395 |
| †16439                   | <i>Dasyscyphella longistipitata</i> Hosoya                              | JAPAN, Kanagawa, Yamakita                              | 2005-04-17        | cupule of <i>Fagus crenata</i>           | 101335                             | AB481239                                 | LC424947 | LC533228 | AB481331 |
| †16527                   | <i>Dasyscyphella montana</i> Raitv.                                     | JAPAN, Nagano, Ueda, Sugadaira Montane Research Center | 2005-05-21        | wood of unidentified-<br>fied tree       | 102336                             | AB481242                                 | LC438577 | LC533241 | AB481336 |
| †16556                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Oita, Kokonoe                                   | 2005-05           | wood of unidenti-<br>fied tree           | 114449                             | UDB0779051                               | LC533153 | LC533257 | LC533198 |
| †16582                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Kanagawa, Yamakita                              | 2005-07-02        | wood of unidenti-<br>fied tree           | 104360                             | AB481249                                 | LC533176 | LC533233 | LC533199 |
| †16606                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Kanagawa, Yamakita                              | 2005-07-03        | wood of unidenti-<br>fied tree           | 114450                             | UDB0779053                               | LC533154 | LC533258 | LC533200 |
| †16609                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Kanagawa, Yamakita                              | 2005-07-03        | wood of <i>Cephalotaxus harringtonia</i> | 101350                             | ††AB705234                               | LC533175 | LC533256 | LC533184 |
| †16639                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Ibaraki, Tsukuba Botanical Garden               | 2006-05-01        | twig of unidentified tree                | 114451                             | UDB0779054                               | LC533155 | LC533259 | LC533201 |
| †25579                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Tokyo, Hongo                                    | 2009-05-25        | twig of unidentified tree                | (FC-1887)                          | UDB0779057                               | LC533146 | LC533250 | LC533191 |
| †32163                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Kanagawa, Odawara                               | 2010-05-14        | twig of unidentified tree                | 114456                             | UDB0779062                               | LC533158 | LC533260 | LC533203 |
| †38452                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Gunma, Naganohara                               | 2013-06-27        | wood of unidenti-<br>fied tree           | 114463                             | ††UDB0779069                             | LC533171 | LC533262 | LC533210 |
| †46416                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | TAIWAN, Taipei                                         | 2012-04-15        | wood of unidenti-<br>fied tree           | (FC-2906)                          | UDB0779067                               | LC533132 | LC533277 | LC549671 |
| †46841                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Gifu, Gujo                                      | 2012-05-28        | wood of unidenti-<br>fied tree           | 114462                             | UDB0779086                               | LC533170 | LC533279 | LC533209 |
| †61773                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Kanagawa, Yokohama                              | 2015-04-01        | twig of unidentified tree                | 114464                             | ††UDB0779074                             | LC533137 | LC533264 | LC533211 |
| †61931                   | <i>Erioscyphella abnormis</i> (Mont.) Baral, Šandová & B. Perić         | JAPAN, Kanagawa, Zushi                                 | 2015-04-16        | wood of unidenti-<br>fied tree           | 114466                             | UDB0779072                               | LC533139 | LC533266 | LC533213 |

| Specimen no.<br>(TNS-F-) | Taxon <sup>1</sup>                                                                                                                          | Collection site                                                            | Collected<br>Date | Host plants and parts                                      | Strain no.<br>(NBRC <sup>2</sup> ) | UNITE/GenBank accession no. <sup>3</sup> |          |          |          |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|-------------------|------------------------------------------------------------|------------------------------------|------------------------------------------|----------|----------|----------|
|                          |                                                                                                                                             |                                                                            |                   |                                                            |                                    | ITS                                      | LSU      | mtSSU    | RPB2     |
| #80478                   | <i>Erioscypbella abnormis</i> (Mont.) Baral, Šandová & B. Perić                                                                             | JAPAN, Shizuoka, Oyama                                                     | 2017-06-26        | twig of unidentified tree                                  | 113934                             | LC424837                                 | LC424949 | LC533283 | LC425009 |
| †26520                   | <i>Erioscypbella boninensis</i> Tochihara & Hosoya                                                                                          | JAPAN, Tokyo, Chichijima Island                                            | 2009-06-28        | trunk of unidentified tree                                 | 114447                             | UDB0779049                               | LC533151 | LC533254 | LC533196 |
| #46419                   | <i>Erioscypbella brasiliensis</i> (Mont.) Baral, Šandová & B. Perić                                                                         | TAIWAN, Taipei                                                             | 2012-04-20        | wood of unidentified tree                                  | (FC-2910)                          | UDB0779068                               | LC533133 | LC533278 | LC549672 |
| #35049                   | <i>Erioscypbella hainanensis</i> (W.Y. Zhuang and Zheng Wang) Hosoya and Tochihara (← <i>Lachnum hainanense</i> W.Y. Zhuang and Zheng Wang) | JAPAN, Niigata, Minamiuonuma                                               | 2010-05-14        | leaf of <i>Quercus glauca</i>                              | 114457                             | UDB0779064                               | LC533168 | LC533274 | LC533205 |
| #35056                   | <i>Erioscypbella hainanensis</i> (W.Y. Zhuang and Zheng Wang) Hosoya and Tochihara (← <i>Lachnum hainanense</i> W.Y. Zhuang and Zheng Wang) | JAPAN, Niigata, Minamiuonuma                                               | 2010-05-14        | leaf of <i>Quercus serrata</i>                             | 114458                             | UDB0779065                               | LC533169 | LC533275 | LC533206 |
| #61775                   | <i>Erioscypbella hainanensis</i> (W.Y. Zhuang and Zheng Wang) Hosoya and Tochihara (← <i>Lachnum hainanense</i> W.Y. Zhuang and Zheng Wang) | JAPAN, Kanagawa, Hiratsuka                                                 | 2015-04-12        | leaf of <i>Quercus myrsinifolia</i>                        | 114465                             | UDB0779071                               | LC533138 | LC533265 | LC533212 |
| #61941                   | <i>Erioscypbella hainanensis</i> (W.Y. Zhuang and Zheng Wang) Hosoya and Tochihara (← <i>Lachnum hainanense</i> W.Y. Zhuang and Zheng Wang) | JAPAN, Kanagawa, Kamakura                                                  | 2015-04-24        | leaf of <i>Quercus glauca</i>                              | 112569                             | UDB0779073                               | LC533140 | LC533280 | LC533214 |
| #65722                   | <i>Erioscypbella hainanensis</i> (W.Y. Zhuang and Zheng Wang) Hosoya and Tochihara (← <i>Lachnum hainanense</i> W.Y. Zhuang and Zheng Wang) | JAPAN, Gunma, Midori                                                       | 2016-04-24        | leaf of <i>Quercus serrata</i> subsp. <i>Mongolicoides</i> | 114469                             | UDB0779076                               | LC533142 | LC533281 | LC533215 |
| #80356                   | <i>Erioscypbella hainanensis</i> (W.Y. Zhuang and Zheng Wang) Hosoya and Tochihara (← <i>Lachnum hainanense</i> W.Y. Zhuang and Zheng Wang) | JAPAN, Kanagawa, Hiratsuka                                                 | 2017-05-18        | leaf of <i>Quercus glauca</i>                              | 114470                             | UDB0779077                               | LC533172 | LC533282 | LC533186 |
| #80371                   | <i>Erioscypbella hainanensis</i> (W.Y. Zhuang and Zheng Wang) Hosoya and Tochihara (← <i>Lachnum hainanense</i> W.Y. Zhuang and Zheng Wang) | JAPAN, Kanagawa, Hiratsuka                                                 | 2017-05-18        | leaf of <i>Castanopsis sieboldii</i>                       | 114472                             | UDB0779078                               | LC533135 | LC533246 | LC533188 |
| †26500                   | <i>Erioscypbella insulae</i> Tochihara & Hosoya                                                                                             | JAPAN, Tokyo, Hahajima Island                                              | 2009-06-24        | wood of unidentified tree                                  | 114445                             | UDB0779060                               | LC533149 | LC533252 | LC533194 |
| #39720                   | <i>Erioscypbella insulae</i> Tochihara & Hosoya                                                                                             | JAPAN, Okinawa, Iriomote Island                                            | 2011-06-12        | bark of unidentified tree                                  | 114459                             | UDB0779063                               | LC533177 | LC533261 | LC533207 |
| #61920                   | <i>Erioscypbella paralushanensis</i> Tochihara & Hosoya                                                                                     | JAPAN, Shizuoka, Arami                                                     | 2015-06-08        | culm of <i>Pleioblastus argenteostriatus</i>               | 114468                             | ††UDB0779075                             | LC533141 | LC533267 | LC533220 |
| †81472                   | <i>Erioscypbella otanii</i> Tochihara                                                                                                       | JAPAN, Hokkaido, Horonobe, Teshio Experimental Forest, Hokkaido University | 2018-07-11        | leaf of <i>Sasa senanensis</i>                             | 114476                             | UDB0779085                               | LC533179 | LC533286 | LC533226 |
| #81272                   | <i>Erioscypbella papillaris</i> Tochihara                                                                                                   | JAPAN, Gunma, Minakami                                                     | 2017-07-16        | leaf of unidentified bamboo                                | 113937                             | UDB0779081                               | LC533161 | LC533285 | LC533204 |
| #80399                   | <i>Erioscypbella sasibrevispora</i> Tochihara & Hosoya                                                                                      | JAPAN, Gunma, Higashi-Agatsuma                                             | 2017-06-06        | sheath of <i>Sasa veitchii</i>                             | —                                  | UDB0779082/<br>LC669470                  | LC533173 | LC533268 | LC533216 |

| Specimen no.<br>(TNS-F.) | Taxon <sup>1</sup>                                                                                                                                             | Collection site                                        | Collected<br>Date | Host plants and parts                                     | Strain no.<br>(NBRC <sup>2</sup> ) | UNITE/GenBank accession no. <sup>3</sup> |          |          |            |
|--------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------|-------------------|-----------------------------------------------------------|------------------------------------|------------------------------------------|----------|----------|------------|
|                          |                                                                                                                                                                |                                                        |                   |                                                           |                                    | ITS                                      | LSU      | mtSSU    | RPB2       |
| ‡81401                   | <i>Erioscyphella sasibrevispora</i> Tochihara & Hosoya                                                                                                         | JAPAN, Hokkaido, Tomakomai                             | 2018-06-16        | culm of <i>Sasa nipponica</i>                             | 114475                             | UDB0779084/<br>LC669472                  | LC533174 | LC533269 | LC533217   |
| ‡26492                   | <i>Erioscyphella sclerotii</i> (A.L. Sm.) Baral, Šandová & B. Perić                                                                                            | JAPAN, Tokyo, Hahajima Island                          | 2009-06-24        | wood of unidentified tree                                 | 114448                             | UDB0779050/<br>LC669438                  | LC533152 | LC533255 | LC533197   |
| ‡38480                   | <i>Erioscyphella sclerotii</i> (A.L. Sm.) Baral, Šandová & B. Perić                                                                                            | TAIWAN, Wulai                                          | 2013-07-12        | twig of unidentified tree                                 | (FC-5208)                          | ‡‡UDB0779070                             | LC533134 | LC533263 | LC549673   |
| ‡16838                   | <i>Erioscyphella sinensis</i> (Z.H. Yu and W.Y. Zhuang) Sasagawa, Tochihara & Hosoya (← <i>Lachnum mapirianum</i> var. <i>sinense</i> Z.H. Yu and W.Y. Zhuang) | JAPAN, Ibaraki, Tsukuba Botanical Garden               | 2007-06-15        | leaf of unidentified broad-leaved tree                    | 104389                             | AB481280                                 | LC533164 | LC533235 | AB481364   |
| ‡80354                   | <i>Erioscyphella sinensis</i> (Z.H. Yu and W.Y. Zhuang) Sasagawa, Tochihara & Hosoya (← <i>Lachnum mapirianum</i> var. <i>sinense</i> Z.H. Yu and W.Y. Zhuang) | JAPAN, Kanagawa, Manazuru                              | 2017-05           | leaf of <i>Castanopsis sieboldii</i>                      | 114471                             | UDB0779083/<br>LC669471                  | LC533143 | LC533245 | LC533187   |
| ‡16841                   | <i>Erioscyphella sinensis</i> (Z.H. Yu and W.Y. Zhuang) Sasagawa, Tochihara & Hosoya (← <i>Lachnum mapirianum</i> var. <i>sinense</i> Z.H. Yu and W.Y. Zhuang) | JAPAN, Ibaraki, Mt. Tsukuba                            | 2007-06-23        | leaf of unidentified broad-leaved tree                    | 104390                             | AB481281                                 | LC533157 | LC533236 | LC533218   |
| ‡32161                   | <i>Erioscyphella sinensis</i> (Z.H. Yu and W.Y. Zhuang) Sasagawa, Tochihara & Hosoya (← <i>Lachnum mapirianum</i> var. <i>sinense</i> Z.H. Yu and W.Y. Zhuang) | JAPAN, Kanagawa, Odawara                               | 2010-05-14        | leaf of <i>Quercus myrsinifolia</i>                       | 113715                             | UDB0779061/<br>LC669449                  | LC533167 | LC533273 | LC533219   |
| ‡16837                   | <i>Erioscyphella sinensis</i> (Z.H. Yu and W.Y. Zhuang) Sasagawa, Tochihara & Hosoya (← <i>Lachnum mapirianum</i> var. <i>sinense</i> Z.H. Yu and W.Y. Zhuang) | JAPAN, Ibaraki, Tsukuba Botanical Garden               | 2007-06-15        | leaf of unidentified broad-leaved tree                    | 114452                             | UDB0779055/<br>LC669443                  | LC533156 | LC533272 | LC533202   |
| ‡81520                   | <i>Incrucipulum ciliare</i> (Schrad.) Baral                                                                                                                    | JAPAN, Shizuoka, Shizuoka                              | 2018-08-18        | leaf of <i>Quercus mongolica</i> subsp. <i>crispula</i>   | 113941                             | LC438566                                 | LC438583 | LC533284 | LC438596   |
| ‡17632                   | <i>Incrucipulum longispineum</i> Sasagawa & Hosoya                                                                                                             | JAPAN, Miyagi, Sendai                                  | 2006-07-29        | leaf of <i>Lyonia ovalifolia</i>                          | 102347                             | AB481256                                 | LC438579 | LC533234 | AB481362   |
| ‡81248                   | <i>Lachnellula calyciformis</i> (Batsch) Dharne                                                                                                                | JAPAN, Hokkaido, Engaru                                | 2017-07-12        | twig of <i>Abies sachalinensis</i>                        | 113935                             | LC438561                                 | LC438574 | LC533247 | LC438590   |
| ‡16529                   | <i>Lachnellula suecica</i> (de Bary ex Fuckel) Nannf.                                                                                                          | JAPAN, Nagano, Ueda, Sugadaira Montane Research Center | 2005-05-21        | twig of <i>Larix kaempferi</i>                            | 101348                             | AB481248                                 | LC424944 | LC533231 | AB481341   |
| ‡16494                   | <i>Lachnum asiaticum</i> (Y. Orani) Raitv.                                                                                                                     | JAPAN, Nagano, Ueda, Sugadaira Montane Research Center | 2005-05-18        | culm of unidentified bamboo                               | 101341                             | AB481251                                 | LC533162 | LC533229 | AB481334   |
| ‡17249                   | <i>Lachnum mapirianum</i> (Pat. & Gaillard) M.P. Sharma                                                                                                        | MALAYSIA, Gerik                                        | 2004-09-07        | leaf of unidentified tree                                 | —                                  | UDB0779088/<br>LC669476                  | LC533182 | —        | LC533223   |
| ‡17245                   | <i>Lachnum mapirianum</i> (Pat. & Gaillard) M.P. Sharma                                                                                                        | MALAYSIA, Gerik                                        | 2004-09-07        | leaf of unidentified tree                                 | —                                  | UDB0779087/<br>LC669475                  | LC533181 | —        | LC533222   |
| ‡16442                   | <i>Lachnum novoguineense</i> var. <i>yunnanicum</i> W.Y. Zhuang                                                                                                | JAPAN, Nagano, Ueda, Sugadaira Montane Research Center | 2005-05-18        | culm of unidentified bamboo                               | 102339                             | AB481270                                 | LC533163 | LC533232 | AB481342   |
| ‡16642                   | <i>Lachnum novoguineense</i> var. <i>yunnanicum</i> W.Y. Zhuang                                                                                                | JAPAN, Ibaraki, Mt. Tsukuba                            | 2006-05-02        | culm of unidentified bamboo                               | 104368                             | AB481271                                 | LC533165 | LC533227 | §§LC533225 |
| ‡11197                   | <i>Lachnum palmae</i> sensu lato (← <i>Lachnum palmae</i> (Kanouse) Spooner)                                                                                   | JAPAN, Shizuoka, Shimoda                               | 2004-07-26        | leaf of <i>Livistona chinensis</i> var. <i>subglobosa</i> | 106495                             | UDB0779047/<br>LC669435                  | LC533166 | LC533248 | LC533185   |

| Specimen no.<br>(TNS-F-) | Taxon <sup>1</sup>                                                               | Collection site                                           | Collected<br>Date | Host plants and parts                                     | Strain no.<br>(NBRC <sup>2</sup> ) | UNITE/GenBank accession no. <sup>3</sup> |          |          |            |
|--------------------------|----------------------------------------------------------------------------------|-----------------------------------------------------------|-------------------|-----------------------------------------------------------|------------------------------------|------------------------------------------|----------|----------|------------|
|                          |                                                                                  |                                                           |                   |                                                           |                                    | ITS                                      | LSU      | mtSSU    | RPB2       |
| ‡13500                   | <i>Lachnum palmarum</i> sensu lato (← <i>Lachnum palmarum</i> (Kanouse) Spooner) | JAPAN, Kagoshima, Yakushima Island                        | 2005-10-19        | leaf of <i>Livistona chinensis</i> var. <i>subglobosa</i> | 114441                             | ††LC425039/<br>UDB779046                 | LC429382 | LC533240 | ‡‡LC431718 |
| ‡17567                   | <i>Lachnum palmarum</i> sensu lato (← <i>Lachnum palmarum</i> (Kanouse) Spooner) | New ZEALAND                                               | 2005-05-28        | leaf of unidentified palm                                 | —                                  | UDB0779089/<br>LC669477                  | LC533183 | LC533288 | —          |
| ‡24588                   | <i>Lachnum palmarum</i> sensu lato (← <i>Lachnum palmarum</i> (Kanouse) Spooner) | JAPAN, Kagoshima, Amami-Oshima                            | 2009-02-24        | leaf of <i>Livistona chinensis</i> var. <i>subglobosa</i> | 114442                             | UDB0779052/<br>LC669440                  | LC533144 | LC533270 | LC533190   |
| ‡24600                   | <i>Lachnum palmarum</i> sensu lato (← <i>Lachnum palmarum</i> (Kanouse) Spooner) | JAPAN, Kagoshima, Amami-Oshima                            | 2009-02-25        | leaf of <i>Livistona chinensis</i> var. <i>subglobosa</i> | 114443                             | UDB0779056/<br>LC669444                  | LC533145 | LC533249 | LC533224   |
| ‡26161                   | <i>Lachnum palmarum</i> sensu lato (← <i>Lachnum palmarum</i> (Kanouse) Spooner) | JAPAN, Tokyo, Chichijima Island                           | 2009-06-27        | leaf of <i>Livistona boninensis</i>                       | 114446                             | UDB0779048/<br>LC669436                  | LC533150 | LC533253 | LC533195   |
| ‡26172                   | <i>Lachnum palmarum</i> sensu lato (← <i>Lachnum palmarum</i> (Kanouse) Spooner) | JAPAN, Tokyo, Kira-Iwojima Island                         | 2009-06-17        | leaf of <i>Livistona chinensis</i> var. <i>subglobosa</i> | (FC-1935)                          | UDB0779058/<br>LC669446                  | LC533147 | LC533251 | LC533192   |
| ‡26185                   | <i>Lachnum palmarum</i> sensu lato (← <i>Lachnum palmarum</i> (Kanouse) Spooner) | JAPAN, Tokyo, Kira-Iwojima Island                         | 2009-06-18        | leaf of <i>Livistona chinensis</i> var. <i>subglobosa</i> | 114444                             | UDB0779059/<br>LC669447                  | LC533148 | LC533271 | LC533193   |
| ‡39729                   | <i>Lachnum palmarum</i> sensu lato (← <i>Lachnum palmarum</i> (Kanouse) Spooner) | JAPAN, Okinawa, Iriomote Island                           | 2011-06-13        | leaf of <i>Livistona chinensis</i> var. <i>subglobosa</i> | 114460                             | UDB0779066/<br>LC669454                  | LC533178 | LC533276 | LC533208   |
| †16501                   | <i>Lachnum pudibundum</i> (Quél.) J. Schröt.                                     | JAPAN, Nagano, Ueda, Sugadaira Montane Research Center    | 2005-05-18        | wood of unidentified tree                                 | 102335                             | AB481259                                 | LC533160 | LC533230 | AB481335   |
| †81229                   | <i>Lachnum rachidicola</i> J.G. Han, Raïtv. & H.D. Shin                          | JAPAN, Hokkaido, Tomakomai, Tomakomai Experimental Forest | 2017-08-09        | petiole of <i>Juglans</i> sp.                             | 114473                             | UDB0779079/<br>LC669467                  | LC533136 | —        | LC533189   |
| †16583                   | <i>Lachnum virgineum</i> (Batsch) P. Karst.                                      | JAPAN, Kanagawa, Yamakita                                 | 2005-07-02        | wood of unidentified tree                                 | 104358                             | AB481268                                 | AB926119 | LC431748 | AB481343   |
| †65625                   | <i>Neodasycephala cerina</i> (Pers.) Spooner                                     | SWITZERLAND, Saïcourt                                     | 2016-06-08        | twig of <i>Crataegus</i> sp.                              | (FC-6068)                          | LC424836                                 | LC424948 | LC533242 | LC425013   |
| †17436                   | <i>Proliferodiscus albiviridis</i> (Sacc.) Spooner                               | JAPAN, Ibaraki, Tsukuba Botanical Garden                  | 2006-07-08        | wood of unidentified tree                                 | 108594                             | LC438558                                 | LC533159 | LC533239 | LC425014   |
| §17909                   | <i>Hyaloscypha spiralis</i> (Velen.) J.G. Han, Hosoya & H.D. Shin                | JAPAN, Kumamoto, Kikuchi                                  | 2005-10-10        | wood of unidentified tree                                 | 108585                             | ††LC438602                               | LC438604 | LC533237 | LC438606   |
| §16472                   | <i>Hymenoscyphus varicosporoides</i> Tubaki                                      | JAPAN, Ibaraki, Kasumigaura                               | 2005-05-05        | wood of unidentified tree                                 | 104355                             | AB926052                                 | LC424952 | LC431746 | AB481329   |
| §18014                   | <i>Urceolella carestiana</i> (Rabenh.) Dennis                                    | JAPAN, Iwate, Hanamaki                                    | 2006-05-23        | stem of <i>Parathelypteris nipponica</i>                  | 108588                             | ††LC438603                               | LC438605 | LC533238 | LC438607   |

† *Lachnaceae* except for *Erioscyphella* and its potential species tentatively identified based on morphology  
‡ *Erioscyphella* or its potential species tentatively identified based on morphology  
§ Outgroup  
| Original taxon name labeled on the specimen is shown enclosed by “(←)” and is only shown when it is different from a name determined in this study.  
¶ Cultures not donated in NBRC is beginning with “FC-”, local suffix in TNS. ‘—’ represents no culture exist and DNA was extracted from apothecia.  
# UNITE accession no. is beginning with ‘UDB’. GenBank accession no. is beginning with ‘AB’ or ‘LC’.  
†† Primer pair ITS1 and ITS4 was used. In ITS sequences without notes (††), primer pair ITS1F and ITS4 was used.  
‡‡ Primer pair RPB2-5F and RPB2-P7R was used.  
§§ Primer pair RPB2-P6Fa and bRPB2-7.1R2 was used.  
|| Primer pair RPB2-P6Fa and RPB2-P7R was used. In RPB2 sequences without any notes (‡‡, §§, ||), primer pair RPB2-P6F and RPB2-P7R was used.




**Figure 1.** ML best-scored phylogenetic tree based on the concatenated dataset of ITS, LSU, mtSSU, and RPB2 constructed using RAxML-NG. MLB/MPBP/BPP are represented on branches in this order. In MLB/MPBP < 50% or BPP < 0.95, a hyphen appears. No evaluation values are shown on branches when MLB/MPBP < 50% and BPP < 0.95. The branch of a clade TNS-F-17245 + 17249 to its most recent common ancestor is only one-third of the intended length due to space limitation.

for LSU, TPM1uf+I+G4 for mtSSU and RPB2 third codon position, GTR+I+G4 for RPB2 first codon position, and TPM3uf+I+G4 for RPB2 second codon position. Sequence matrix containing missing data typically yield multiple trees residing on a phylogenetic terrace (Sanderson et al. 2011; Biczok et al. 2018). Therefore, we checked if the best-scored-tree did not lie on a terrace using the Python tool called ‘terrapihy’ implemented in RAxML-NG 0.9.0.

MP analysis was conducted using PAUP\* 4.0a 167 (Swofford 2002). All substitutions were treated as unordered and of equal weights. All gaps were treated as missing data. A heuristic parsimony search was carried out with 1,000 replicates of random step addition, with a tree bisection reconnection (TBR) branch swapping algorithm, Multrees option on, Steepest descent modification option on, and branch collapse option set to MinBrLen. Bootstrap values (MPBP; Felsenstein 1985) were estimated from 1,000 replicates of heuristic searches, with random taxon addition, TBR branch swapping, and Multrees options off.

BI analysis was based on MrBayes 3.2.7a (Ronquist et al. 2012) under the substitution model SYM+I+G4 for ITS, GTR+I+G4 for LSU and RPB2 first codon positions, HKY+I+G4 for mtSSU and RPB2 third codon positions, and F81+I for RPB2 second codon position. Two separate Metropolis-Coupled Markov Chains of Monte Carlo (MCMCMC) ran simultaneously starting from random trees for 20 million generations, and trees were sampled every 500 generations. The average standard deviation of split frequencies (ASDSF) and effective sample size (ESS) were checked using Tracer 1.7.1 (Rambaut 2018a) as an indication of convergence. Using post-burn-in trees, a 50% majority rule consensus tree was generated, and Bayesian posterior probabilities (BPP) were calculated to evaluate node supports. Trees were visualized using FigTree 1.4.4 (Rambaut 2018b) based on the ML, MP, and BI analyses respectively. Branches with MLBP and MPBP > 90% and BPP > 0.95 were regarded as strongly supported.

ITS-based species delimitation analyses (Fig. 2)

To maximize the number of ITS sequences, we used the UNITE Species Hypotheses (SH) system provided by the UNITE database (Kõljalg et al. 2013; Nilsson et al. 2015; GBIF 2018; Kõljalg et al. 2020). In the UNITE SH system, all fungal ITS sequences are periodically divided into species-level clusters (species hypothesis; SH) at optional sequence-distance thresholds (0%–3% in 0.5% steps), each of which is assigned to a unique UNITE SH code represented by a digital objective identifier (DOI) accessible from internet (Kõljalg et al. 2016, 2020; Nilsson et al. 2015).

Based on the UNITE SH system, we collected ITS sequences of *Erioscyphella* in the following process: a) selectivity of closely related sequences: for every ITS sequence newly obtained from TNS specimens (= query sequences, 49 sequences), UNITE SH code at the 3% threshold value were searched in the UNITE database to gather sequences in wider scope, and all sequences within the UNITE SH code were downloaded. b) selectivity based on taxon names: using the UNITE search page, ITS sequences named *Erioscyphella* were searched, because only closely related sequences to query sequences are filtered under the a) criterion. Sequences with synonyms of *Erioscyphella* species were also searched, because the UNITE lookup function is not supported by any backbone taxonomies to integrate synonyms. Sequences satisfying criterion a) or




**Figure 2.** Diagrammatic representation showing the species delimitation analyses using ITS sequences.

b) were downloaded for ITS-based species recognition. The obtained ITS sequences were clustered into SHs based on an OTU clustering method, hierarchical clustering method, and two coalescent-based methods. For all ITS sequences, ITS1, 5.8S, and ITS2 regions were extracted using ITSx (Nilsson et al. 2010) to construct an accurate ITS dataset, because the inclusion of segments of adjacent regions (such as a small subunit of 18S rRNA or LSU) may decrease the accuracy of the calculation of ITS distances (Nilsson et al. 2010). OTU clustering was executed using VSEARCH v2.17.2 (Rognes et al. 2016) implemented in the Qiime 2 microbiome analysis platform (Bolyen et al. 2019).

The concatenated dataset of extracted ITS1, 5.8S, and ITS2 was incorporated into VSEARCH, and OTU clustering at 97% and 98.5% similarity thresholds were performed using the ‘-cluster\_fast’ option. Hierarchical clustering based on pairwise sequence distances was executed using the Assemble Species by Automatic Partitioning (ASAP) method (Puillandre et al. 2021). The datasets of extracted ITS1, 5.8S, and ITS2 were separately aligned using MAFFT 7 under the Q-INS-i option and edited using trimAl v1.2 (Capella-Gutiérrez et al. 2009) under the ‘-gappyout’ option. The concatenated dataset of the three aligned partitions was analyzed using ASAP web (<https://bioinfo.mnhn.fr/abi/public/asap/asapweb.html>). Jukes-Cantor (JC69) was selected as a substitution model for computing pairwise distances of sequences. As phylogeny-based species delimitation methods, the generalized mixed Yule-coalescent (GMYC) model (Pons et al. 2006; Fujisawa and Barraclough 2013) and the Poisson Tree Processes (PTP) model (Zhang et al. 2013) were used. In both models, specia-

tion (species-level differentiation) and coalescence (population-level differentiation) are identified based on the length of phylogenetic trees. GMYC requires the use of phylogenetic trees following the molecular clock model (= ultrametric tree) because it detects transition points from speciation to coalescence focusing on the time axis, while PTP does not require ultrametric tree as it focuses on the number of nucleotide substitutions. Ultrametric trees were estimated using BEAST v2.6.3. (Bouckaert et al. 2019). The ITS dataset was divided into ITS1, 5.8S, and ITS2, and suitable substitution models GTR+G for ITS1 and JC+G for 5.8S and ITS2 estimated using Modeltest-NG 0.1.6. were applied. To estimate branch length, a Yule model and a relaxed clock with a log-normal distribution were selected. MCMC chains were run for  $1.5 \times 10^8$  generations and sampled every 1,000 generations. After each run, convergence was checked using Tracer 1.7.1, and the first 10% were discarded as burn-in. A consensus tree was generated using TreeAnnotator v1.10.4 in BEAST package, from 150,000 generated trees except for the first 10% regarded as burn-in. A single-threshold species delimitation analysis based on GMYC was conducted using the R package 'splits' (Fujisawa and Barraclough 2013).

For the species delimitation analyses using PTP, an unrooted ML phylogenetic tree was constructed using RAxML-NG 0.9.0. The analysis used ITS1, 5.8S, and ITS2 partitions, aligned as previously described, under the substitution models TIM2+G4 for ITS1, TPM2+I+G4 for 5.8S, and GTR+I+G4 for ITS2, estimated using Modeltest-NG 0.1.6. based on the AIC. The species delimitation analysis was executed using the generated ML best-scored tree with the bPTP web server (<https://species.h-its.org/>). The MCMC run was set to 500,000 generations and burn-in rate was set to 0.1. The convergence of MCMC runs was visually checked. In ML and Bayesian results, a result generating fewer SHs was adopted to avoid excessive species division.

SHs generated in the species delimitation analyses and the UNITE SHs at 3% and 1.5% threshold values were compared with one another.

## Species recognition

In the present study, we initially recognized species boundaries based on the two criteria:

1. Forming a monophyletic group in the phylogenetic analyses based on multi-gene data (Fig. 1).
2. Members can be distinguished based on morphological and/or common ecological features (such as host plants).

Species boundaries recognized by 1. and 2. were cross-checked based on the results of ITS-based species delimitation analyses. When the species boundaries are supported by the majority (= more than four methods) of the seven species delimitation methods (UNITE SH at 3% threshold, UNITE SH at 1.5% threshold, VSEARCH 97% similarity, VSEARCH 98.5% similarity, ASAP, GMYC, and PTP) (Fig. 3), we regard the species as reasonable and carry out taxonomic treatments if necessary.




**Figure 3.** Species delimitation analyses using ITS sequences of *Erioscyphella* and its potential members. Clusters based on UNITE SH at 3% and 1.5% threshold values at UNITE v8.2, VSEARCH at 97% and 98.5% threshold values, ASAP, GMYC, and PTP are displayed. Schematic phylogenetic relationships are shown using the ultrametric tree constructed for the GMYC analysis. The taxon names shown on the tree branches follow the results of the present study.

## Results

### Taxon sampling from TNS specimens

Forty-nine specimens in TNS were identified as candidates of *Erioscyphella* and morphologically identified as *E. abnormis*, *E. brasiliensis*, *E. sclerotii*, *Lachnum hainanense* W.Y. Zhuang & Zheng Wang, *L. mapirianum* (Pat. & Gaillard) M.P. Sharma, *Lachnum mapirianum* var. *sinense* Z.H. Yu, W.Y. Zhuang, *Lachnum novoguineense* var. *yun-nanicum* W.Y. Zhuang, and *L. palmae* (Kanouse) Spooner (Table 1), together with six species of *Erioscyphella* described here as new (*E. boninensis*, *E. insulae*, *E. otanii*, *E. papillaris*, *E. paralushanensis*, and *E. sasibrevispora*], Table 1).

### Phylogenetic analyses

The molecular phylogenetic analyses were based on 70 specimens selected from TNS (Table 1). The concatenated sequence matrix was composed of 2488 bp (sites 1–332 for ITS, 333–1108 for LSU, 1109–1828 for mtSSU, and 1829–2488 for RPB2). In the matrix, the following parts were treated as missing data: TNS-F-17245, 17249, and 81229 for mtSSU, and TNS-F-17567 for RPB2. The matrix was registered in TreeBase (<http://purl.org/phylo/treebase/phyloids/study/TB2:S28477>).

Among the four ML trees based on each region, no conflicts were found in clades with support > 70% (Suppl. material 1: Fig. S1). Therefore, we considered these four regions to be combinable, and phylogenetic analyses were based on the concatenated sequence matrix. In the ML analysis, the best-scored tree generated did not reside on the phylogenetic terrace. In the MP analysis, 766 nucleotide substitution sites were detected, 601 of which were parsimony-informative. A total of 182,630 equally parsimonious trees were generated with tree length = 2,985 steps, consistency index (CI) = 0.38, retention index (RI) = 0.73, and rescaled consistency index (RC) = 0.28. In the BI analysis, when two runs reached 20 million generations and the first 10,000 trees (25%) of generated trees were excluded, ASDSF was observed to fall below 0.004 and ESS of all parameters was over 200. The first 10,000 trees were discarded as burn-in. A 50% majority rule consensus tree was constructed and BPP was calculated based on the remaining 30,000 trees.

As no topological contradictions occurred among the ML best-scored tree, MP 50% majority-rule consensus tree, and BI 50% majority-rule consensus tree, only ML tree was illustrated, and MLBS, MPBS, and BPP were plotted on its branches (Fig. 1).

Based on the phylogenetic analyses, 49 candidates of *Erioscyphella* formed a strongly supported clade (= Clade A, MLBP = 100%/MPBP = 100%/BPP = 1.00), apart from the clade of *Lachnum* sensu stricto (= *L. asiaticum* (Y. Otani) Raitv., *L. pudibundum* (Quél.) J. Schröt., *L. rachidicola* J.G. Han, Raitv. & H.D. Shin, and *L. virgineum* (Batsch) P. Karst.) [type of *Lachnum*]) (Fig. 1). Clade A and *Proliferodiscus alboviridis* formed a relatively strongly supported clade (Clade B, MLBP = 78%, MPBP = 82%, BPP = 1.00).

Within Clade A, each morphologically identified species and variety formed strongly supported monophyletic groups of their own (Fig. 1), and five strongly supported

subclades were recognized (Clade I–V, Fig. 1). *Lachnum mapirianum* (TNS-F-17545, 17249) and *E. insulae* (TNS-F-26500, 39720) did not belong to any subclade. Clade I was composed of *E. boninensis*, *E. paralushanensis*, *L. hainanense*, and *L. mapirianum* var. *sinense*. Within Clade I, only *E. paralushanensis* occurred on bamboo sheaths, while others occurred on fallen leaves of broad-leaved trees. Clade II was composed only of *L. palmae*, which occurred on the palm petioles. Clade III was composed of *E. otanii* and *E. papillaris* occurring on bamboo leaves. Clade IV was composed of *L. novoguineense* var. *yunnanicum*, and *E. sasibrevispora*, occurring on bamboo sheaths. Clade V was composed of *E. abnormis*, *E. brasiliensis*, and *E. sclerotii*, occurring on wood.

## Morphological characters within Clade A

Members of Clade A had totally and densely granulate, hyaline to brown, thin-walled hairs, fusiform to long filiform ascospores, ectal excipulum composed of *textura prismatica* to *textura angularis*, asci lacking croziers at the bases, and smooth walled ectal excipulum cells. Exceptionally, *E. sasibrevispora*, *L. hainanense* (Hosoya et al. 2013), and *L. novoguineense* var. *yunnanicum* W.Y. Zhuang had croziers and *E. boninensis* had granulated ectal excipulum.

Moreover, hairs of Clade A lacked crystals, but were equipped with apical amorphous materials and/or resinous materials. In the present study, “crystals” refers to amber colored materials that positioned near the hair apices and were regular-shaped (e.g. tetrahedral materials, masses of needle-like materials, or cross-shaped materials), described by Raitviir (2002), Suková (2005) or Tochihara and Hosoya (2019). “Resinous materials” refers to colored, refractive, irregular-shaped materials attached on any parts of hairs, described by Spooner (1987). Crystals and resinous materials are easily detached from hairs and broken into fragments in the squash mount. “Apical amorphous materials” is termed uniquely in this study, and refers to hyaline to brown, refractive, irregular-shaped materials positioned outside the hair apices. They are usually small and inconspicuous cap-like shaped, and conspicuously globular in some species. Apical amorphous materials do not grow to big masses and are not easily detached from hairs in the squash mount.

In Clade A, members except for *E. boninensis*, *E. sasibrevispora* and *L. novoguineense* var. *yunnanicum* had apical amorphous materials, and *E. boninensis*, *E. paralushanensis*, and *L. palmae* complex also had resinous materials (see figures of described species and Suppl. material 1: Fig. S2).

## ITS-based species delimitation analyses

In UNITE v8.3, 87 ITS sequences were clustered into 23 SHs at 3% and 26 SHs at 1.5% threshold values (Table 2, Fig. 3). The UNITE SH code for each SH is presented in Table 2. In OTU clustering using VSEARCH, 87 ITS sequences were clustered into 25 SHs at 97% similarity and 28 SHs at 98.5% similarity (Table 2, Fig. 3). VSEARCH SH codes (allocated in this study uniquely; VSH97\_1 to VSH97\_25, VSH985\_1 to VSH985\_28) are shown in Table 2.

Table 2. ITS sequences analyzed by the species delimitation analyses.

| ITS sequence<br>GenBank/UNITE<br>accession no.   | TNS-F<br>speci-<br>men<br>no. | Reference (initial<br>appearance) | Taxon name<br>(ultimately<br>allocated in<br>this study) | UNITE taxon<br>name    | INSDC taxon<br>name    | Country                      | Host plants and<br>parts                       | UNITE SH code<br>(DOI) at 3%<br>threshold | UNITE SH code<br>(DOI) at 1.5%<br>threshold | VSEARCH<br>SH at 97%<br>similarity | VSEARCH<br>SH at 98.5%<br>similarity |
|--------------------------------------------------|-------------------------------|-----------------------------------|----------------------------------------------------------|------------------------|------------------------|------------------------------|------------------------------------------------|-------------------------------------------|---------------------------------------------|------------------------------------|--------------------------------------|
| AB267634                                         |                               | Miyoshi et al. (2007)             | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | JAPAN, Ehime                 | twig of <i>Citrus junos</i>                    | SH1155612.08FU                            | SH1522994.08FU                              | VSH97_1                            | VSH985_2                             |
| AB267636 (dupli-<br>cate; AB267635)              |                               | Miyoshi et al. (2007)             | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | JAPAN, Ehime                 | twig of <i>Citrus junos</i>                    | SH1155612.08FU                            | SH1522994.08FU                              | VSH97_1                            | VSH985_2                             |
| AB267641 (dupli-<br>cate; AB267639,<br>AB267640) |                               | Miyoshi et al. (2007)             | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | JAPAN, Tokushima             | twig of <i>Citrus junos</i>                    | SH1155612.08FU                            | SH1522994.08FU                              | VSH97_1                            | VSH985_2                             |
| AB267642                                         |                               | Miyoshi et al. (2007)             | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | JAPAN, Tokushima             | twig of <i>Citrus junos</i>                    | SH1155612.08FU                            | SH1522994.08FU                              | VSH97_1                            | VSH985_2                             |
| JF937578                                         |                               | Zhao and Zhuang<br>(2011)         | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | CHINA                        | (unspecified)                                  | SH1155612.08FU                            | SH1522994.08FU                              | VSH97_1                            | VSH985_2                             |
| JN033395                                         |                               | Han et al. (2014)                 | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | KOREA                        | Wood                                           | SH1155612.08FU                            | SH1522994.08FU                              | VSH97_1                            | VSH985_2                             |
| UDB0779067/<br>LC669455                          | 46416                         | this study                        | <i>E. abnormis</i>                                       | -                      | -                      | TAIWAN, Taipei               | wood of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1522994.08FU                              | VSH97_1                            | VSH985_2                             |
| UDB0779074/<br>LC669462                          | 61773                         | this study                        | <i>E. abnormis</i>                                       | -                      | -                      | JAPAN, Kanagawa,<br>Yokohama | twig of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1522994.08FU                              | VSH97_1                            | VSH985_2                             |
| MK584950                                         |                               | Ekanayaka et al.<br>(2019)        | <i>E. abnormis</i>                                       | <i>E. abnormis</i>     | <i>E. abnormis</i>     | CHINA, Yunnan                | (unspecified)                                  | SH1155612.08FU                            | †SH1522994.08FU                             | VSH97_1                            | VSH985_2                             |
| AB267637                                         |                               | Miyoshi et al. (2007)             | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | JAPAN, Nara                  | Twig                                           | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_1                             |
| AB267638                                         |                               | Miyoshi et al. (2007)             | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | JAPAN, Shizuoka              | Twig                                           | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_1                             |
| AB481249                                         | 16582                         | Hosoya et al. (2010)              | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | JAPAN, Kanagawa,<br>Yamakita | wood of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_1                            | VSH985_1                             |
| AB705234                                         | 16609                         | Zhao et al. (2012)                | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | JAPAN, Kanagawa,<br>Yamakita | wood of <i>Cephalo-<br/>taxus harringtonia</i> | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_1                            | VSH985_1                             |
| LC424837                                         | 80478                         | this study                        | <i>E. abnormis</i>                                       | -                      | -                      | JAPAN, Shizuoka,<br>Oyama    | twig of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_3                             |
| MG712307                                         |                               | unpublished                       | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | CHINA                        | (unspecified)                                  | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_3                             |
| MK282241                                         |                               | unpublished                       | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | (unspecified)                | (unspecified)                                  | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_1                             |
| MK584957                                         |                               | Ekanayaka et al.<br>(2019)        | <i>E. abnormis</i>                                       | <i>E. aseptata</i>     | <i>E. aseptata</i>     | THAILAND,<br>Chiang Rai      | (unspecified)                                  | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_1                             |
| MN082536                                         |                               | unpublished                       | <i>E. abnormis</i>                                       | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i> | (unspecified)                | (unspecified)                                  | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_1                            | VSH985_1                             |
| MT995055                                         |                               | unpublished                       | <i>E. abnormis</i><br>(misregis-<br>tered?)              | <i>Chapsa patens</i>   | <i>Chapsa patens</i>   | (unspecified)                | (unspecified)                                  | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_1                            | VSH985_1                             |
| MW007918                                         |                               | unpublished                       | <i>E. abnormis</i><br>(misregis-<br>tered?)              | <i>Chapsa patens</i>   | <i>Chapsa patens</i>   | (unspecified)                | (unspecified)                                  | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_3                             |

| ITS sequence<br>GenBank/UNITE<br>accession no. | TNS-F<br>speci-<br>men<br>no. | Reference (initial<br>appearance) | Taxon name<br>(ultimately<br>allocated in<br>this study)                              | UNITE taxon<br>name    | INSDC taxon<br>name                                      | Country                                        | Host plants and<br>parts                       | UNITE SH code<br>(DOI) at 3%<br>threshold | UNITE SH code<br>(DOI) at 1.5%<br>threshold | VSEARCH<br>SH at 97%<br>similarity | VSEARCH<br>SH at 98.5%<br>similarity |
|------------------------------------------------|-------------------------------|-----------------------------------|---------------------------------------------------------------------------------------|------------------------|----------------------------------------------------------|------------------------------------------------|------------------------------------------------|-------------------------------------------|---------------------------------------------|------------------------------------|--------------------------------------|
| UDB0779051/<br>LC669439                        | 16556                         | this study                        | <i>E. abnormis</i>                                                                    | -                      | -                                                        | JAPAN, Oita,<br>Kokonoe                        | wood of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_1                             |
| UDB0779053/<br>LC669441                        | 16606                         | this study                        | <i>E. abnormis</i>                                                                    | -                      | -                                                        | JAPAN, Kanagawa,<br>Yamakita                   | wood of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_3                             |
| UDB0779054/<br>LC669442                        | 16639                         | this study                        | <i>E. abnormis</i>                                                                    | -                      | -                                                        | JAPAN, Ibaraki,<br>Tsukuba Botanical<br>Garden | twig of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_3                             |
| UDB0779057/<br>LC669445                        | 25579                         | this study                        | <i>E. abnormis</i>                                                                    | -                      | -                                                        | JAPAN, Tokyo,<br>Hongo                         | twig of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_3                             |
| UDB0779062/<br>LC669450                        | 32163                         | this study                        | <i>E. abnormis</i>                                                                    | -                      | -                                                        | JAPAN, Kanagawa,<br>Odawara                    | twig of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_3                             |
| UDB0779069/<br>LC669457                        | 38452                         | this study                        | <i>E. abnormis</i>                                                                    | -                      | -                                                        | JAPAN, Gunma,<br>Naganohara                    | twig of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_1                            | VSH985_1                             |
| UDB0779072/<br>LC669460                        | 61931                         | this study                        | <i>E. abnormis</i>                                                                    | -                      | -                                                        | JAPAN, Kanagawa,<br>Zushi                      | twig of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_2                            | VSH985_3                             |
| UDB0779086/<br>LC669474                        | 46841                         | this study                        | <i>E. abnormis</i>                                                                    | -                      | -                                                        | JAPAN, Gifu, Gujo                              | twig of unidenti-<br>fied tree                 | SH1155612.08FU                            | SH1523013.08FU                              | VSH97_1                            | VSH985_1                             |
| AB481250                                       | 16617                         | Hosoya et al. (2010)              | <i>E. abnormis</i>                                                                    | <i>Lachnum abnorme</i> | <i>Lachnum abnorme</i>                                   | JAPAN, Kanagawa,<br>Yamakita                   | twig of unidenti-<br>fied tree                 | ‡SH1155612.08FU                           | ‡SH1523013.08FU                             | VSH97_1                            | VSH985_1                             |
| UDB0779055/<br>LC669443                        | 16837                         | this study                        | <i>E. sinensis</i><br>(← <i>Lachnum</i><br><i>mapirianum</i><br>var. <i>sinense</i> ) | -                      | -                                                        | JAPAN, Ibaraki,<br>Tsukuba Botanical<br>Garden | leaf of unidentified<br>broad-leaved tree      | SH1155682.08FU                            | SH1523107.08FU                              | VSH97_4                            | VSH985_5                             |
| AB481280                                       | 16838                         | Hosoya et al. (2010)              | <i>E. sinensis</i><br>(← <i>Lachnum</i><br><i>mapirianum</i><br>var. <i>sinense</i> ) | <i>Lachnum</i> sp.     | <i>Lachnum</i> ( <i>Lach-</i><br><i>num</i> sp. FC-2355) | JAPAN, Ibaraki,<br>Tsukuba Botanical<br>Garden | leaf of unidentified<br>broad-leaved tree      | SH1155682.08FU                            | SH1523107.08FU                              | VSH97_4                            | VSH985_5                             |
| AB481281                                       | 16841                         | Hosoya et al. (2010)              | <i>E. sinensis</i><br>(← <i>Lachnum</i><br><i>mapirianum</i><br>var. <i>sinense</i> ) | <i>Lachnum</i> sp.     | <i>Lachnum</i> ( <i>Lach-</i><br><i>num</i> sp. FC-2358) | JAPAN, Ibaraki,<br>Mt. Tsukuba                 | leaf of unidentified<br>broad-leaved tree      | SH1155682.08FU                            | SH1523107.08FU                              | VSH97_4                            | VSH985_5                             |
| UDB0779061/<br>LC669449                        | 32161                         | this study                        | <i>E. sinensis</i><br>(← <i>Lachnum</i><br><i>mapirianum</i><br>var. <i>sinense</i> ) | -                      | -                                                        | JAPAN, Kanagawa,<br>Odawara                    | leaf of <i>Quercus</i><br><i>myrsinifolia</i>  | SH1155682.08FU                            | SH1523107.08FU                              | VSH97_4                            | VSH985_5                             |
| UDB0779083/<br>LC669471                        | 80354                         | this study                        | <i>E. sinensis</i><br>(← <i>Lachnum</i><br><i>mapirianum</i><br>var. <i>sinense</i> ) | -                      | -                                                        | JAPAN, Kanagawa,<br>Manazuru                   | leaf of <i>Castanopsis</i><br><i>sieboldii</i> | ‡SH1155682.08FU                           | ‡SH1523107.08FU                             | VSH97_4                            | VSH985_5                             |

| ITS sequence<br>GenBank/UNITE<br>accession no. | TNS-F<br>speci-<br>men<br>no. | Reference (initial<br>appearance) | Taxon name<br>(ultimately<br>allocated in<br>this study)          | UNITE taxon<br>name              | INSDC taxon<br>name              | Country                           | Host plants and<br>parts                                                                       | UNITE SH code<br>(DOI) at 3%<br>threshold | UNITE SH code<br>(DOI) at 1.5%<br>threshold | VSEARCH<br>SH at 97%<br>similarity | VSEARCH<br>SH at 98.5%<br>similarity |
|------------------------------------------------|-------------------------------|-----------------------------------|-------------------------------------------------------------------|----------------------------------|----------------------------------|-----------------------------------|------------------------------------------------------------------------------------------------|-------------------------------------------|---------------------------------------------|------------------------------------|--------------------------------------|
| UDB023346                                      |                               | unpublished                       | <i>E. curvispora</i>                                              | <i>E. curvispora</i>             | -                                | MONTENEGRO,<br>Žijevero Mountains | needle of <i>Pinus<br/>heldreichii</i>                                                         | SH1155703.08FU                            | SH1523136.08FU                              | VSH97_12                           | VSH985_14                            |
| MH190414                                       |                               | Perić and Baral<br>(2014)         | <i>E. curvispora</i>                                              | <i>E. curvispora</i>             | <i>E. curvispora</i>             | MONTENEGRO,<br>Žijevero Mountains | needle of <i>Pinus<br/>heldreichii</i>                                                         | †SH1155703.08FU                           | †SH1523136.08FU                             | VSH97_12                           | VSH985_14                            |
| JF937580                                       |                               | Zhao and Zhuang<br>(2011)         | <i>E. brasiliensis</i>                                            | <i>Lachnum brasil-<br/>iense</i> | <i>Lachnum brasil-<br/>iense</i> | CHINA                             | (unspecified)                                                                                  | SH1155705.08FU                            | SH1523142.08FU                              | VSH97_6                            | VSH985_7                             |
| MK584953                                       |                               | Ekanayaka et al.<br>(2019)        | <i>E. brasiliensis</i>                                            | <i>E. brasiliensis</i>           | <i>E. brasiliensis</i>           | (unspecified)                     | (unspecified)                                                                                  | SH1155705.08FU                            | SH1523142.08FU                              | VSH97_6                            | VSH985_7                             |
| MK584967                                       |                               | Ekanayaka et al.<br>(2019)        | <i>E. brasiliensis</i>                                            | <i>E. brasiliensis</i>           | <i>E. brasiliensis</i>           | THAILAND,<br>Chiang Rai           | (unspecified)                                                                                  | SH1155705.08FU                            | SH1523142.08FU                              | VSH97_6                            | VSH985_7                             |
| UDB0779068/<br>LC669456                        | 46419                         | this study                        | <i>E. brasiliensis</i>                                            | -                                | -                                | TAIWAN, Taipei                    | wood of unidenti-<br>fied tree                                                                 | SH1155705.08FU                            | SH1523142.08FU                              | VSH97_6                            | VSH985_7                             |
| JF937579                                       |                               | Zhao and Zhuang<br>(2011)         | <i>E. brasiliensis</i>                                            | <i>Lachnum brasil-<br/>iense</i> | <i>Lachnum brasil-<br/>iense</i> | CHINA                             | (unspecified)                                                                                  | †SH1155705.08FU                           | †SH1523142.08FU                             | VSH97_6                            | VSH985_7                             |
| KX501132                                       |                               | Tello and Baral<br>(2016)         | <i>E. lunata</i>                                                  | <i>E. lunata</i>                 | <i>E. lunata</i>                 | SPAIN, Andalucía                  | needle of <i>Pinus<br/>nigra</i> subsp. <i>nigra</i>                                           | †SH1155760.08FU                           | †SH1523257.08FU                             | VSH97_18                           | VSH985_19                            |
| JX984680                                       |                               | unpublished                       | <i>E. hai-<br/>nanensis</i><br>(← <i>Lachnum<br/>hainanense</i> ) | Hyaloscyphaceae                  | Fungi (uncultured<br>fungus)     | KOREA, Seoul                      | (Total suspended<br>particulate matter<br>(TSP) in urban air<br>during non-Asian<br>dust days) | SH1155844.08FU                            | SH1523423.08FU                              | VSH97_3                            | VSH985_4                             |
| UDB0779064/<br>LC669452                        | 35049                         | this study                        | <i>E. hai-<br/>nanensis</i><br>(← <i>Lachnum<br/>hainanense</i> ) | -                                | -                                | JAPAN, Niigata,<br>Minamiuonuma   | leaf of <i>Quercus<br/>glauca</i>                                                              | SH1155844.08FU                            | SH1523423.08FU                              | VSH97_3                            | VSH985_4                             |
| UDB0779065/<br>LC669453                        | 35056                         | this study                        | <i>E. hai-<br/>nanensis</i><br>(← <i>Lachnum<br/>hainanense</i> ) | -                                | -                                | JAPAN, Niigata,<br>Minamiuonuma   | leaf of <i>Quercus<br/>serrata</i>                                                             | SH1155844.08FU                            | SH1523423.08FU                              | VSH97_3                            | VSH985_4                             |
| UDB0779073/<br>LC669461                        | 61941                         | this study                        | <i>E. hai-<br/>nanensis</i><br>(← <i>Lachnum<br/>hainanense</i> ) | -                                | -                                | JAPAN, Kanagawa,<br>Kamakura      | leaf of <i>Quercus<br/>glauca</i>                                                              | SH1155844.08FU                            | SH1523423.08FU                              | VSH97_3                            | VSH985_4                             |
| UDB0779076/<br>LC669464                        | 65722                         | this study                        | <i>E. hai-<br/>nanensis</i><br>(← <i>Lachnum<br/>hainanense</i> ) | -                                | -                                | JAPAN, Gunma,<br>Midori           | leaf of <i>Quercus<br/>serrata</i> subsp.<br><i>mongolicoides</i>                              | SH1155844.08FU                            | SH1523423.08FU                              | VSH97_3                            | VSH985_4                             |

| ITS sequence<br>GenBank/UNITE<br>accession no. | TNS-F<br>speci-<br>men<br>no. | Reference (initial<br>appearance) | Taxon name<br>(ultimately<br>allocated in<br>this study) | UNITE taxon<br>name      | INSDC taxon<br>name                                 | Country                                                         | Host plants and<br>parts             | UNITE SH code<br>(DOI) at 3%<br>threshold | UNITE SH code<br>(DOI) at 1.5%<br>threshold | VSEARCH<br>SH at 97%<br>similarity | VSEARCH<br>SH at 98.5%<br>similarity |
|------------------------------------------------|-------------------------------|-----------------------------------|----------------------------------------------------------|--------------------------|-----------------------------------------------------|-----------------------------------------------------------------|--------------------------------------|-------------------------------------------|---------------------------------------------|------------------------------------|--------------------------------------|
| MK282242                                       |                               | unpublished                       | <i>E. hai-nanensis</i><br>(← <i>Lachnum hainanense</i> ) | <i>Lachnum</i> sp.       | <i>Lachnum albidulum</i>                            | KOREA                                                           | (unspecified)                        | SH1155844.08FU                            | †SH1523423.08FU                             | VSH97_3                            | VSH985_4                             |
| UDB0779077/<br>LC669465                        | 80356                         | this study                        | <i>E. hai-nanensis</i><br>(← <i>Lachnum hainanense</i> ) | -                        | -                                                   | JAPAN, Kanagawa,<br>Hiratsuka                                   | leaf of <i>Quercus glauca</i>        | SH1155844.08FU                            | SH3597461.08FU                              | VSH97_3                            | VSH985_9                             |
| UDB0779078/<br>LC669466                        | 80371                         | this study                        | <i>E. hai-nanensis</i><br>(← <i>Lachnum hainanense</i> ) | -                        | -                                                   | JAPAN, Kanagawa,<br>Hiratsuka                                   | leaf of <i>Castanopsis sieboldii</i> | SH1155844.08FU                            | SH3597461.08FU                              | VSH97_3                            | VSH985_9                             |
| UDB0779071/<br>LC669459                        | 61775                         | this study                        | <i>E. hai-nanensis</i><br>(← <i>Lachnum hainanense</i> ) | -                        | -                                                   | JAPAN, Kanagawa,<br>Hiratsuka                                   | leaf of <i>Quercus myrsinifolia</i>  | †SH1155844.08FU                           | †SH3597461.08FU                             | VSH97_3                            | VSH985_9                             |
| UDB0779050/<br>LC669438                        | 26492                         | this study                        | <i>E. sclerotii</i>                                      | -                        | -                                                   | JAPAN, Tokyo,<br>Habajima Island                                | wood of unidentified tree            | SH1155848.08FU                            | SH1523429.08FU                              | VSH97_5                            | VSH985_6                             |
| JF937584                                       |                               | Zhao and Zhuang<br>(2011)         | <i>E. sclerotii</i>                                      | <i>Lachnum sclerotii</i> | <i>Lachnum sclerotii</i>                            | CHINA                                                           | (unspecified)                        | SH1155848.08FU                            | SH1523429.08FU                              | VSH97_5                            | VSH985_6                             |
| MK584951                                       |                               | Ekanayaka et al.<br>(2019)        | <i>E. sclerotii</i>                                      | <i>E. sclerotii</i>      | <i>E. sclerotii</i>                                 | THAILAND,<br>Chiang Rai                                         | (unspecified)                        | SH1155848.08FU                            | SH1523429.08FU                              | VSH97_5                            | VSH985_6                             |
| UDB0779070/<br>LC669458                        | 38480                         | this study                        | <i>E. sclerotii</i>                                      | -                        | -                                                   | TAIWAN, Wulai                                                   | twig of unidentified tree            | SH1155848.08FU                            | SH1523429.08FU                              | VSH97_5                            | VSH985_6                             |
| MK584969                                       |                               | Ekanayaka et al.<br>(2019)        | <i>E. sclerotii</i>                                      | <i>E. sclerotii</i>      | <i>E. sclerotii</i>                                 | THAILAND,<br>Chiang Rai                                         | (unspecified)                        | †SH1155848.08FU                           | †SH1523429.08FU                             | VSH97_5                            | VSH985_6                             |
| AB481271                                       | 16642                         | Hosoya et al. (2010)              | <i>Lachnum novoguineense</i><br>var. <i>yunnanicum</i>   | <i>Lachnum</i> sp.       | <i>Lachnum</i> sp.<br>( <i>Lachnum</i> sp. FC-2211) | JAPAN, Ibaraki,<br>Mt. Tsukuba                                  | culm of unidentified bamboo          | SH1236904.08FU                            | SH1648536.08FU                              | VSH97_10                           | VSH985_12                            |
| AB481270                                       | 16442                         | Hosoya et al. (2010)              | <i>Lachnum novoguineense</i><br>var. <i>yunnanicum</i>   | <i>Lachnum</i> sp.       | <i>Lachnum</i> sp.<br>( <i>Lachnum</i> sp. FC-2117) | JAPAN, Nagano,<br>Ueda, Sugadaira<br>Montane Research<br>Center | culm of unidentified bamboo          | †SH1236904.08FU                           | †SH1648536.08FU                             | VSH97_10                           | VSH985_12                            |
| MK584965                                       |                               | Ekanayaka et al.<br>(2019)        | <i>E. alba</i>                                           | <i>E. alba</i>           | <i>E. alba</i>                                      | THAILAND,<br>Chiang Mai                                         | (unspecified)                        | †SH2596405.08FU                           | †SH2712425.08FU                             | VSH97_22                           | VSH985_25                            |
| AB267647                                       |                               | Miyoshi et al. (2007)             | <i>Lachnum palmæ</i> sensu lato                          | <i>Lachnum palmæ</i>     | <i>Lachnum palmæ</i>                                | JAPAN, Oita                                                     | leaf of <i>Livistona chinensis</i>   | SH1149764.08FU                            | SH1515235.08FU                              | VSH97_7                            | VSH985_8                             |

| ITS sequence<br>GenBank/UNITE<br>accession no. | TNS-F<br>speci-<br>men<br>no. | Reference (initial<br>appearance) | Taxon name<br>(ultimately<br>allocated in<br>this study) | UNITE taxon<br>name      | INSDC taxon<br>name      | Country                                    | Host plants and<br>parts                                            | UNITE SH code<br>(DOI) at 3%<br>threshold | UNITE SH code<br>(DOI) at 1.5%<br>threshold | VSEARCH<br>SH at 97%<br>similarity | VSEARCH<br>SH at 98.5%<br>similarity |
|------------------------------------------------|-------------------------------|-----------------------------------|----------------------------------------------------------|--------------------------|--------------------------|--------------------------------------------|---------------------------------------------------------------------|-------------------------------------------|---------------------------------------------|------------------------------------|--------------------------------------|
| LC425039<br>(duplicate;<br>UDB0779046)         | 13500                         | Johnston et al.<br>(2019)         | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | <i>Lachnum palmae</i>    | <i>Lachnum palmae</i>    | JAPAN, Kagoshi-<br>ma, Yakushima<br>Island | leaf of <i>Livistona<br/>chinensis</i> var. <i>sub-<br/>globosa</i> | SH1149764.08FU                            | SH1515235.08FU                              | VSH97_7                            | VSH985_8                             |
| UDB0779066/<br>LC669454                        | 39729                         | this study                        | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | -                        | -                        | JAPAN, Okinawa,<br>Iriomote Island         | leaf of <i>Livistona<br/>chinensis</i> var. <i>sub-<br/>globosa</i> | SH1149764.08FU                            | SH1515235.08FU                              | VSH97_7                            | VSH985_8                             |
| MG283320                                       |                               | Zhao et al. (2018)                | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | <i>Lachnum palmae</i>    | <i>Lachnum palmae</i>    | CHINA, Linzhou                             | root of <i>Przewalskia<br/>tangutica</i> (endo-<br>phyte)           | †SH1149764.08FU                           | †SH1515235.08FU                             | VSH97_7                            | VSH985_8                             |
| UDB0779089/<br>LC669477                        | 17567                         | this study                        | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | -                        | -                        | NEW ZEALAND                                | leaf of unidentified<br>palm                                        | SH2594271.08FU                            | SH2709065.08FU                              | VSH97_15                           | VSH985_16                            |
| MH921862                                       |                               | unpublished                       | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | <i>Lachnum palmae</i>    | <i>Lachnum palmae</i>    | NEW ZEALAND                                | unidentified part of<br><i>Rhopalostylis sapida</i>                 | †SH2594271.08FU                           | †SH2709065.08FU                             | VSH97_15                           | VSH985_16                            |
| UDB0779052/<br>LC669440                        | 24588                         | this study                        | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | -                        | -                        | JAPAN, Ka-<br>goshima, Amami-<br>Oshima    | leaf of <i>Livistona<br/>chinensis</i> var. <i>sub-<br/>globosa</i> | SH3569651.08FU                            | SH3597456.08FU                              | VSH97_9                            | VSH985_17                            |
| UDB0779047/<br>LC669435                        | 11197                         | this study                        | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | -                        | -                        | JAPAN, Shizuoka,<br>Shimoda                | leaf of <i>Livistona<br/>chinensis</i> var. <i>sub-<br/>globosa</i> | †SH3569651.08FU                           | †SH3597456.08FU                             | VSH97_9                            | VSH985_17                            |
| UDB0779048/<br>LC669436                        | 26161                         | this study                        | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | -                        | -                        | JAPAN, Tokyo,<br>Chichijima Island         | leaf of <i>Livistona<br/>boninensis</i>                             | SH3569651.08FU                            | SH3597457.08FU                              | VSH97_9                            | VSH985_11                            |
| UDB0779058/<br>LC669446                        | 26172                         | this study                        | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | -                        | -                        | JAPAN, Tokyo,<br>Kita-Iwojima Island       | leaf of <i>Livistona<br/>chinensis</i> var. <i>sub-<br/>globosa</i> | SH3569651.08FU                            | SH3597457.08FU                              | VSH97_16                           | VSH985_11                            |
| UDB0779059/<br>LC669447                        | 26185                         | this study                        | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | -                        | -                        | JAPAN, Tokyo,<br>Kita-Iwojima Island       | leaf of <i>Livistona<br/>chinensis</i> var. <i>sub-<br/>globosa</i> | SH3569651.08FU                            | †SH3597457.08FU                             | VSH97_16                           | VSH985_11                            |
| UDB0779056/<br>LC669444                        | 24600                         | this study                        | <i>Lachnum<br/>palmae</i> sensu<br>lato                  | -                        | -                        | JAPAN, Ka-<br>goshima, Amami-<br>Oshima    | leaf of <i>Livistona<br/>chinensis</i> var. <i>sub-<br/>globosa</i> | †SH3569653.08FU                           | †SH3597459.08FU                             | VSH97_25                           | VSH985_28                            |
| U58640                                         |                               | Cantrell and Hanlin<br>(1997)     | <i>E. euterpes</i>                                       | <i>Lachnum euterpes</i>  | <i>Lachnum euterpes</i>  | PUERTO RICO                                | (unspecified)                                                       | †SH1236906.08FU                           | †SH1648538.08FU                             | VSH97_21                           | VSH985_24                            |
| KT384413                                       |                               | Ekanayaka et al.<br>(2019)        | <i>E. fusiformis</i>                                     | <i>Lachnum fusiforme</i> | <i>Lachnum fusiforme</i> | THAILAND                                   | dead stems                                                          | †SH1236907.08FU                           | †SH1648539.08FU                             | VSH97_11                           | VSH985_13                            |
| MK584948                                       |                               | Ekanayaka et al.<br>(2019)        | <i>E. fusiformis</i>                                     | <i>Lachnum fusiforme</i> | <i>Lachnum fusiforme</i> | CHINA                                      | dead stems                                                          | SH1236907.08FU                            | SH1648539.08FU                              | VSH97_11                           | VSH985_13                            |

| ITS sequence<br>GenBank/UNITE<br>accession no. | TNS-F<br>speci-<br>men<br>no. | Reference (initial<br>appearance) | Taxon name<br>(ultimately<br>allocated in<br>this study) | UNITE taxon<br>name             | INSDC taxon<br>name             | Country                                                                                | Host plants and<br>parts                                        | UNITE SH code<br>(DOI) at 3%<br>threshold | UNITE SH code<br>(DOI) at 1.5%<br>threshold | VSEARCH<br>SH at 97%<br>similarity | VSEARCH<br>SH at 98.5%<br>similarity |
|------------------------------------------------|-------------------------------|-----------------------------------|----------------------------------------------------------|---------------------------------|---------------------------------|----------------------------------------------------------------------------------------|-----------------------------------------------------------------|-------------------------------------------|---------------------------------------------|------------------------------------|--------------------------------------|
| UDB0779049/<br>LC669437                        | 26520                         | this study                        | <i>E. boninensis</i>                                     | -                               | -                               | JAPAN, Tokyo,<br>Hahajima Island                                                       | wood of unidenti-<br>fied tree                                  | †SH3569652.08FU                           | †SH3597458.08FU                             | VSH97_20                           | VSH985_21                            |
| UDB0779060/<br>LC669448                        | 26500                         | this study                        | <i>E. insulae</i>                                        | -                               | -                               | JAPAN, Tokyo,<br>Hahajima Island                                                       | wood of unidenti-<br>fied tree                                  | SH3569654.08FU                            | SH3597460.08FU                              | VSH97_14                           | VSH985_15                            |
| UDB0779063/<br>LC669451                        | 39720                         | this study                        | <i>E. insulae</i>                                        | -                               | -                               | JAPAN, Okinawa,<br>Iriomote Island                                                     | bark of unidenti-<br>fied tree                                  | †SH3569654.08FU                           | †SH3597460.08FU                             | VSH97_14                           | VSH985_15                            |
| UDB0779075/<br>LC669463                        | 61920                         | this study                        | <i>E. paralusha-<br/>nensis</i>                          | -                               | -                               | JAPAN, Shizuoka,<br>Arami                                                              | culm of <i>Pleioblastus<br/>argenteostriatus</i>                | †SH3569655.08FU                           | †SH3597462.08FU                             | VSH97_19                           | VSH985_20                            |
| AF505515                                       |                               |                                   | <i>E. lushanensis</i>                                    | <i>Lachnum lusha-<br/>nense</i> | <i>Lachnum lusha-<br/>nense</i> | (unspecified)                                                                          | (unspecified)                                                   | †SH1155706.08FU                           | †SH1523143.08FU                             | VSH97_8                            | VSH985_10                            |
| JF937582                                       |                               | Zhao and Zhuang<br>(2011)         | <i>E. lushanensis</i>                                    | <i>Lachnum lusha-<br/>nense</i> | <i>Lachnum lusha-<br/>nense</i> | CHINA                                                                                  | (unspecified)                                                   | SH1155706.08FU                            | SH1523143.08FU                              | VSH97_8                            | VSH985_10                            |
| MG434782                                       |                               | unpublished                       | <i>E. lushanensis</i>                                    | <i>Erioscyphella</i> sp.        | <i>E. lushanensis</i>           | INDIA, Tangmarg                                                                        | root tips of <i>Pinus<br/>wallichiana</i> (ecto-<br>mycorrhiza) | (unassigned)                              | (unassigned)                                | VSH97_8                            | VSH985_10                            |
| UDB0779081/<br>LC669469                        | 81272                         | this study                        | <i>E. papillaris</i>                                     | -                               | -                               | JAPAN, Gunma,<br>Minakami                                                              | leaf of unidentified<br>bamboo                                  | †SH3569656.08FU                           | †SH3597463.08FU                             | VSH97_23                           | VSH985_26                            |
| UDB0779084/<br>LC669472                        | 81401                         | this study                        | <i>E. sasibrevis-<br/>pora</i>                           | -                               | -                               | JAPAN, Hokkaido,<br>Tomakomai                                                          | culm of <i>Sasa nip-<br/>ponica</i>                             | SH3569657.08FU                            | SH3597464.08FU                              | VSH97_13                           | VSH985_23                            |
| UDB0779082/<br>LC669470                        | 80399                         | this study                        | <i>E. sasibrevis-<br/>pora</i>                           | -                               | -                               | JAPAN, Gunma,<br>Higashi-Agatsuma                                                      | sheath of <i>Sasa<br/>vetitchii</i>                             | †SH3569657.08FU                           | †SH3597464.08FU                             | VSH97_13                           | VSH985_22                            |
| UDB0779085/<br>LC669473                        | 81472                         | this study                        | <i>E. otanii</i>                                         | -                               | -                               | JAPAN, Hokkaido,<br>Horonobe, Teshio<br>Experimental<br>Forest, Hokkaido<br>University | leaf of <i>Sasa sena-<br/>nensis</i>                            | †SH3569658.08FU                           | †SH3597465.08FU                             | VSH97_24                           | VSH985_27                            |
| UDB0779087/<br>LC669475                        | 17245                         | this study                        | <i>Lachnum<br/>mapirianum</i>                            | -                               | -                               | MALAYSIA, Gerik                                                                        | leaf of unidenti-<br>fied tree                                  | †SH3569659.08FU                           | †SH3597466.08FU                             | VSH97_17                           | VSH985_18                            |
| UDB0779088/<br>LC669476                        | 17249                         | this study                        | <i>Lachnum<br/>mapirianum</i>                            | -                               | -                               | MALAYSIA, Gerik                                                                        | leaf of unidenti-<br>fied tree                                  | SH3569659.08FU                            | SH3597466.08FU                              | VSH97_17                           | VSH985_18                            |

† Representative sequence of each SH

‡ Reference sequence of each SH

The extracted and aligned ITS sequences were composed of three partitions, ITS1 (162 bp), 5.8S (157 bp), and ITS2 (142 bp). The concatenated ITS sequence matrix was registered in TreeBase (<http://purl.org/phylo/treebase/phyloids/study/TB2:S28473>). In the ASAP analysis, the concatenated dataset of these partitions (461 bp) was input, and 87 ITS sequences were clustered into 18 SHs with the lowest asap-score, reflecting better partitioning (Suppl. material 1: Fig. S3). In the GMYC analysis, 29 SHs were delimited (Suppl. material 1: Fig. S4). The ultrametric tree constructed for the GMYC analysis is available in TreeBase (<http://purl.org/phylo/treebase/phyloids/study/TB2:S28473>). For the PTP analyses, an ML best-scored tree was constructed (Suppl. material 1: Fig. S5). PTP analyses delimited 23 SHs in the Bayesian support and 26 SHs in the ML support (Suppl. material 1 Fig. S6), and the former was adopted.

Comparing the number of SHs generated by different clustering methods and applied thresholds, 18 SHs by ASAP, and 23 SHs by UNITE SH at 3% threshold represented the lowest SH numbers (Fig. 3; Table 2). The ASAP results were too rough to delimit the boundaries of *E. abnormis*, *E. boninensis*, *E. brasiliensis*, *E. curvispora*, and *E. sclerotii*. SH-classification recognized by UNITE SH at 3% threshold mostly corresponded to taxon names originally assigned to sequences.

Comparing the results of seven species delimitation methods (UNITE SH at 3% threshold, UNITE SH at 1.5% threshold, VSEARCH 97% similarity, VSEARCH 98.5% similarity, ASAP, GMYC, and PTP), sequences labeled as *E. alba*, *E. brasiliensis*, *E. curvispora*, *E. euterpes*, *E. fusiformis*, *E. lunata*, *E. sclerotii*, *L. mapirianum*, *L. mapirianum* var. *sinense*, *L. novoguineense* var. *yunnanica*, and six new species candidates were distinguished as separate clusters by more than four delimitation methods (Fig. 3). These species clusters did not contradict with morphological/ecological and phylogenetic relationships (Fig. 1). Seven sequences labeled as *L. hainanense* were clustered into one SH by four species delimitation analyses, and part of the SHs included a sequence labeled as *Lachnum albidulum* (Fig. 3).

*Erioscyphella abnormis*, *E. aseptate*, and *L. palmae* did not form separate clusters supported by majority of four species delimitation analyses (Fig. 3). Sequences labeled as *E. abnormis* were clustered into one to four SHs, and some SHs included sequences labeled as *Chapsa patens* (Nyl.) Frisch, *E. aseptata*, *E. brasiliensis*, and *E. sclerotii* (Fig. 3). Twelve sequences labeled as *L. palmae* were clustered into four to six SHs (Fig. 3).

## Discussion

### Generic delimitation and generic concept of *Erioscyphella*

We accepted Clade A as a monophyletic unit for *Erioscyphella* which is supported by morphology. Although Clade B comprised Clade A together with *P. alboviridis*, Clade B should not be regarded as a genus delimitation of *Erioscyphella*, because *Proliferodiscus* differs from members of Clade A in having apothecia proliferating from the margins continuously and thick-walled and coarsely warted hairs (Haines and Dumont 1983; Spooner 1987). All members of Clade A are distinguishable from the other

lachnaceous genera. In contrast to *Erioscyphella*, *Albotricha* and *Dasyscyphella* are distinguished by hair apices with no granulation (Hosoya et al. 2010), *Brunnipila*, *Capitotricha*, and *Incrucipulum* by hair-crystals (Baral and Krieglsteiner 1985; Tochiara and Hosoya 2019), and *Lachnellula* by ectal excipulum composed of *textura globosa* to *textura oblita* (Dharne 1965). Typical members of Clade A can be easily segregated from *Neodasyscypha*, because the characteristic features of *Neodasyscypha*, such as dark-brown hairs, ectal-excipulum structure, and ellipsoid to fusoid ascospores < 10 µm long (Spooner 1987), are rare in Clade A. Among members of Clade A and *Lachnum* sensu stricto, the shape and length of ascospores were continuous (Fig. 4), as indicated by Haines and Dumont (1984). However, ascospores longer than 15–20 µm were restricted to Clade A (Fig. 4). Moreover, most members of Clade A have hairs with apical amorphous materials, which are not seen in *Lachnum* sensu stricto. Members of Clade A usually also have hairs not swelling at the apices and distantly septate, as Perić and Baral (2014) pointed out for three tropical members, while members of *Lachnum* have swelling apices. The combination of such characters allows us to differentiate typical members of *Erioscyphella* from *Lachnum*.

In summary, *Erioscyphella* is still difficult to define solely based on morphology because of multiple exceptional characters continuous to other genera, but its typical members could be recognizable mainly by the hair structures and ascospore length. Based on members of Clade A, *Erioscyphella* is tentatively described as follows: apothecia occurring on dead hardwood leaves, rotten wood, bamboo sheaths, bamboo leaves or palm leaves; asci mostly arising from simple septa, but occasionally from croziers; ascospores fusiform to long needle-shaped, aseptate to multi-septate; paraphyses filiform to narrowly lanceolate, shortly exceeding the asci, but rarely lanceolate and long exceeding the asci; hairs straight or irregularly curved, usually not swollen at the apices, thin-walled, hyaline, but sometimes brown, totally and densely granulated, usually distantly septate, without needle-like or three-dimensional shaped crystals but mostly equipped with hyaline to brown apical amorphous materials, and/or resinous materials at any part of hairs; walls of ectal excipulum cells smooth but granulate in one species.

Perić and Baral (2014) pointed out that “yellow hymenium derived from carotenoid” is one of the common characters of *Erioscyphella*. This feature was not discussed in this study because some specimens were not observed when fresh; the hymenium color is variable (usually white hymenium becomes yellow) between fresh and dried states in lachnaceous species.

### Host selectivity of *Erioscyphella*

In *Erioscyphella*, the tendency of selectivity of species to host plants or parts occurs across the genus. Each subclade within *Erioscyphella* (Clade I–V) generally shared tendencies toward host selectivity as follows: Clade I on leaves of broad-leaved trees, except for *E. paralushanensis* occurring on bamboo sheaths, Clade II on palm leaves, Clade III on bamboo leaves, Clade IV on bamboo sheaths, and Clade V on rotten wood (Fig. 1). The results showed that selectivity to host plants, and parts of *Erioscyphella*, was acquired as apomorphic characters during speciation.




**Figure 4.** Comparison of ascospores of Clade A (= *Erioscyphella*) and the clade of *Lachnum* sensu stricto in Fig. 1. Subclade numbers for members of Clade A in Fig. 1 are shown in parentheses. Bars show variation of ascospore length within each species.

### Is *Erioscyphella* limited to ‘tropical’ zones?

*Erioscyphella* (long-spored *Lachnum*) has long been known as the tropical genus in Lachnaceae (Dennis 1954; Spooner 1987; Guatimosim et al. 2016). Most long-spored species were described from tropical areas of Latin America (Dennis 1954) and tropical to temperate areas of Australasia (Spooner 1987). However, the new species or new combinations proposed in this study were reported from Japan in subtropical areas (*E. boninensis* and *E. insulae*), temperate area (*E. hainanensis*, *E. paralushanensis*, and *E. sinensis*) and cool-temperate to subarctic areas (*E. otanii*, *E. papillaris*, and *E. sasibrevispora*), showing that *Erioscyphella* is not limited to tropical zones, but is also distributed in temperate to subarctic zones in the northern hemisphere.

### Ascal iodine reactions seen in *E. papillaris*

Iodine reactions of the ascus apical apparatus have been classified into several types (inamyloid, hemiamyloid [Type RB and RR, and euamyloid Type BB]) (Baral 2009), and the reaction ‘MLZ- without KOH pretreatment and MLZ+ with KOH pretreatment’, observed in *E. papillaris* (Fig. 11E1 and Fig. E2) has been restricted to the type of hemiamyloid. However, the apical apparatus of *E. papillaris* showed a dark blue reaction in IKI without KOH pretreatment (Fig. 11E3), while the hemiamyloid apparatus usually shows a red reaction under these conditions. The hemiamyloid ascus apparatus could show IKI-blue without KOH pretreatment due to long storage in the herbarium (Baral 2009), but this is not applicable for the material of *E. papillaris*, which has been maintained for only two years in herbarium until observed. Therefore, we assessed the iodine reaction of *E. papillaris* as a new type, and color reactions with various solutions of the species should be further examined using new materials, because there are few apothecia in the type specimen.

### Species-level taxonomic treatment of *Erioscyphella*

In this study, we carried out taxonomic treatment for species which were distinguished by morphology/ecology and phylogenetic analyses, and formed single clusters in species delimitation analyses. Based on this criteria, six undescribed species of *Erioscyphella* have been proposed as new species of *Erioscyphella* [*E. boninensis*, *E. insulae*, *E. otanii*, *E. papillaris*, *E. paralushanensis*, and *E. sasibrevispora*], and *Lachnum hainanense* and *L. mapirianum* var. *sinense* have been proposed as new members of *Erioscyphella*. Interpretation of species boundaries of *L. hainanense* was discussed in the taxonomy chapter. For new species and new combinations, Japanese names were also denominated for wider use of Japanese mycologists or amateurs.

In the phylogenetic analyses, Malaysian materials of *L. mapirianum* (TNS-F-17245, 17249) and Japanese materials of *L. novoguineense* var. *yunnanicum* (TNS-F-16442, 16642) were also found to be members of *Erioscyphella* (Fig. 1). However, we hesitate to transfer the two species into *Erioscyphella*, as we cannot guarantee the identification accuracy of the materials, because of inadequate type information of the two species.

Taxonomic assessments of *E. abnormis*, *L. aseptata*, and *L. palmae*, which were not accepted as independent species in species delimitation analyses, are discussed below.

### Taxonomy of *E. abnormis* and its related species

In the species delimitation analyses, sequences labeled as *E. abnormis* formed a single SH at UNITE SH 3% threshold (DOI: SH1155612.08FU) and divided into two to four SHs at UNITE SH 1.5% threshold, VSEARCH, and GMYC (Fig. 3).

In ASAP, sequences labeled as *E. abnormis* belong to a single SH, but the SH also contained sequences labeled as *Chapsa patens*, *E. aseptata*, *E. brasiliensis*, *E. curvispora*,

and *E. sclerotii* (Fig. 3). However, the phylogenetic analyses revealed that *E. brasiliensis*, and *E. sclerotii* are separate from the clade of *E. abnormis* (Fig. 1), suggesting that the two species are different from *E. abnormis*. Although *E. curvispora* was not included in the phylogenetic analyses (Fig. 1), the apparent morphological and ecological differentiation (Perić and Baral 2014) and low similarity of ITS (< 97%) with members of *E. abnormis* (Fig. 3) suggest that *E. curvispora* is different from *E. abnormis*.

*Erioscyphella aseptata* was originally described in Thailand and characterized by having aseptate ascospores, unlike *E. abnormis* or *E. sclerotii* with septate ascospores (Ekanayaka et al. 2019). However, the species delimitation analyses in this study suggested the difficulty of delimiting *E. aseptata* (MK584957) from *E. abnormis* (Fig. 3), suggesting that *E. aseptata* is a morphologically atypical (aseptate-ascospored) individual of *E. abnormis*.

Although two ITS sequences of *C. patens* (MT995055 = specimen no. FJ19131 and MW007918 = specimen no. FJ19049) were positioned in SHs dominated by *E. abnormis*, LSU and mtSSU sequences of FJ19131 and LSU sequence of FJ19049 were closely related to *Chapsa* spp. [Graphidaceae, Ostropales]. Since Lachnaceae and Graphidaceae are phylogenetically distant, the two ITS sequences MT995055 and MW007918 have been misidentified.

Considering that the monophyly of *E. abnormis* is strongly supported (Fig. 1) and members of the species share high ITS similarities (> 97%, compiled into SH1155612.08FU) (Fig. 3, Table 2), *E. abnormis* is accepted here as a species with some intraspecific morphological and phylogenetic variation.

## Taxonomy of ‘*Lachnum*’ *palmae*

*Lachnum palmae* formed a strongly supported clade in the phylogenetic analyses (Clade II in Fig. 1). They also shared strong selectivity to palm leaves and characteristic morphology such as thick-walled asci, hairs with resinous materials and apical amorphous materials (Suppl. material 1: Fig. S2) and ectal excipulum composed of thick-walled prismatic cells and interwoven hyphae. However, sequences labeled as *L. palmae* were divided into 4 to 7 SHs in all species delimitation analyses (Fig. 3), indicating that *L. palmae* is a species complex that includes multiple potential sister species. At present, we avoid creating new species from the complex, because the morphological and ecological differences detected among SHs are not enough to delimit species boundaries, although the size of asci and ascospores differ among some SHs, as shown in Fig. 4. Phylogenetic analyses revealed that members of the *L. palmae* complex belonged to *Erioscyphella* (Fig. 1). However, we could not judge which SH within the complex is equivalent to *L. palmae* as originally described from Honduras by Kanouse (1941) and redescribed by Spooner (1987) from the type plus another specimen from New Zealand. There are no *L. palmae* sequences from the tropical American type locality, so phylogenetic characterization and recombination of the species were avoided in the present study.

## Taxonomy

### *Erioscyphella boninensis* Tochihara & Hosoya, sp. nov.

MycoBank No: 835702

Figs 5, 6

**Diagnosis.** Differs from all other *Erioscyphella* species by the granulate walls of the ectal excipular cells.

**Holotype.** JAPAN, Bonin Islands, Chichijima Island, Mt. Tsutsujiyama, 27.060556, 142.222500, ca 270 m, 28 Jun. 2009, on fallen leaves of *Pittosporum boninense*, T.Hosoya (TNS-F-26520).

**GenBank/UNITE no. ex holotype.** LC669437/UDB0779049 (ITS), LC533151 (LSU), LC533254 (mtSSU), LC533196 (RPB2).

**Etymology.** Referring to the type locality Bonin Islands.

**Japanese name.** Ogasawara-cha-hina-no-chawantake.

**Description.** Apothecia scattered, superficial, 0.5–1.0 mm in diameter, having well-developed stipes, up to 1.5 mm high, cream to pale brown, externally covered with short and shiny hairs. Disc concave, cream to pale yellow. Ectal excipulum *textura prismatica* composed of long elongated cells to *textura angularis*, 6–25 × 5–13 µm, hyaline to relatively brown colored, somewhat thick-walled; cell walls covered by granules with a similar appearance to those on hairs. Stipe composed of *textura prismatica* with a granulate surface as ectal excipular cells. Medullary excipulum *textura intricata* of hyaline hyphae up to 3 µm wide. Hairs straight, cylindrical, 38–62 × 2.5–4.0 µm, hyaline, completely covered by brown granules, 2–3-septate, thin-walled, arising from swelling cells completely covered by granules; apex lacking crystals or apical amorphous materials, equipped with amber-colored resinous materials dissolvable with CB/LA at a little below the apex. Asci (36–)37.7–44(–46) × (3.5–)3.6–4.2(–4.5) µm (av. 41 ± 3.2 × 3.9 ± 0.3 µm, n = 16), 8-spored, cylindrical-clavate; pore blue in MLZ without 3% KOH pretreatment; croziers absent at the basal septa. Ascospores (9–)10–12.3(–13) × 1.2–1.7(–1.8) µm (av. 11 ± 1.2 × 1.5 ± 0.2 µm, n = 16), Q = (6.3–)6.9–9.2(–10) (av. 7.8 ± 1.5, n = 16), fusiform, aseptate. Paraphyses straight, up to 2.5 µm wide, septate, exceeding the asci up to 5 µm, narrowly lanceolate.

**Culture characteristics.** Colony of NBRC 114447/TNS-F-26520 on PDA umbonate forming a dome-shape, slightly sulcate. Context not shiny, velvety, buff at the center, paler toward the margin, dark buff from the reverse. Sectors and zonation absent. Aerial mycelium white or buff, dense cottony, forming white mycelium strands except in the margin. Margin distinct, entire, flat. Asexual morph absent.

**Distribution.** JAPAN. (Bonin Islands). Known only from the type locality.

**Notes.** Granulation on the surface of the ectal excipular cells has been observed only in *Incrucipulum* in Lachnaceae (Baral and Krieglsteiner 1985; Tochihara and Hosoya 2019), and *E. boninensis* is the first report for such a character in *Erioscyphella*




**Figure 5.** *Erioscyphella boninensis* TNS-F-26520 (Holotype) **A** dried apothecia **B** pure culture on PDA (NBRC 114447) **C** ascus **D** ascus pore MLZ (+) **E** ascospores **F** paraphyses **G** ectal excipular cells **H** ectal excipular cells with red granules **I** hairs with resinous matters arising from ectal excipular cells. Mounted in CB/LA (**C**, **E–I**), MLZ (**D**). Scale bars: 1 mm (**A**); 10 µm (**C–I**).

(Fig. 5H, 6E). Phylogenetic analysis revealed that *E. boninensis* is closely related to *E. paralushanensis* (Fig. 1). The two species (Clade IA, Fig. 1) have colored granules on hairs and forming red mycelia on PDA. However, granulation of ectal excipulum is seen only in *E. boninensis*.

***Erioscyphella hainanensis* (W.Y. Zhuang and Zheng Wang) Hosoya and Tochihara, comb. nov.**

MycoBank No: 835707

≡ *Lachnum hainanense* W.Y. Zhuang & Zheng Wang, Mycotaxon 67: 25 (1998).

**Diagnosis.** Forming apothecia with long stipes and long hairs. Differing *E. sinensis* in much shorter ascospores.

**Japanese name.** Shii-Kashi-hina-no-chawantake.

**Specimens examined.** JAPAN, Niigata, Minamiuonuma, 37.056808, 138.80705, ca 720 m, 14 May 2010, on fallen leaves of *Quercus glauca*, T.Hosoya




**Figure 6.** *Erioscyphella boninensis* TNS-F-26520 (Holotype) **A** ascospores **B** apothecium **C** vertical section of an apothecium **D** expansion of a vertical section of an apothecium **E** ectal excipular cells **F** asci **G** paraphyses **H** hairs.

(TNS-F-35049). Ibid (TNS-F-35056). JAPAN, Kanagawa, Hiratsuka, 35.33861111, 139.285, ca 80 m, 12 Apr. 2015, on fallen leaves of *Q. myrsinifolia*, M.Nakajima (TNS-F-61775). JAPAN, Kanagawa, Kamakura, 35.30756, 139.51958, ca 40 m, 24 Apr. 2015, on fallen leaves of *Q. serrata*, M.Nakajima (TNS-F-61941). JAPAN, Gunma, Midori, 36.476684, 139.242771, ca 510 m, 9 May 2016, on fallen leaves of *Q. serrata*, K.Furuya (TNS-F-65722). JAPAN, Kanagawa, Hiratsuka, 35.340139, 139.287167, ca 60 m, 18 May 2017, on fallen leaves of *Q. glauca*, Y.Tochiara (TNS-F-80356). The same locality, on fallen leaves of *Castanopsis sieboldii*, Y. Tochiara (TNS-F-80371).

**Distribution.** CHINA (Hainan), JAPAN (Honshu: Kanto region).

**Notes.** Based on the UNITE SH system at a 3% threshold, ITS sequences of this species were integrated into a single SH (DOI: SH1155844.08FU). SH1155844.08FU included sequences labeled as ‘*Hyaloscyphaceae*’ (JX984680) in UNITE and ‘*L. albidulum*’ (MK282242) in INSDC (Table 2). JX984680 was sequenced from air samples in Seoul, South Korea, and was not tied to any fungal specimens or cultures. *Lachnum albidulum* is common on leathery dicot leaves of the old and new world tropics (Haines 1992). *Erioscyphella hainanensis* resembles *L. albidulum* in morphology, but *L. albidulum* has yellow resinous substances at the tip of apothecial hairs and occurs on dead leaves of Rubiaceae (Haines 1992), whereas *E. hainanensis* lacks resinous sub-

stances and occurs on leaves of broad-leaved trees (Zhuang and Wang 1998b; Hosoya et al. 2013). Therefore, we presume that MK282242, coexisting with *L. hainanense* in every SH, was misidentified as *L. albidulum*. No sequences are available for *L. albidulum* specimens from the type locality. *Lachnum hainanense* was therefore judged as acceptable species, and recombined into *Erioscyphella*.

*Erioscyphella hainanensis* resembles *E. sinensis* in occurring on dead leaves of *Quercus* spp. or *Castanopsis* spp. However, *E. hainanensis* has much shorter ascospores than *E. sinensis*. In this study, presence of minute, hyaline apical amorphous materials and absence of any crystals or resinous materials were confirmed in both species (Suppl. material 1: Fig. S2).

***Erioscyphella insulae* Tochihara & Hosoya, sp. nov.**

MycoBank No: 835703

Figs 7, 8

**Diagnosis.** Characterized by pure white apothecia unlike related species *Lachnum nothofagi*, and two-layered ectal excipulum.

**Holotype.** JAPAN, Okinawa, Yaeyama, Taketomi, Iriomote Island, Otomi, 24.297458, 123.866128, ca 50 m, 12 Jun. 2011, on fallen bark of unidentified tree, T.Fukiharu (TNS-F-39720).

**GenBank/UNITE no. ex holotype.** LC669451/UDB0779063 (ITS), LC533177 (LSU), LC533261 (mtSSU), LC533207 (RPB2).

**Other specimens examined.** JAPAN, Bonin Islands, Hahajima Island, Sekimon, 26.666686, 142.152222, ca 260 m, 24 Jun. 2009, on fallen bark of unidentified tree, T.Hosoya (TNS-F-26485, 26500).

**Etymology.** Referring to the occurrence of the species on remote islands in Japan.

**Japanese name.** Shima-hina-no-chawantake.

**Description.** Apothecia gregarious, superficial, 0.7–1.4(–2.5) mm in diameter, short- and thick-stipitate, up to 0.8 mm high, externally white to cream throughout but sometimes pale brown in the lower parts, covered with white hairs. Disc concave, cream to pale yellow (fresh state not observed). Ectal excipulum composed of two layers: outer layer *textura angularis*, up to 20 µm thick, 3–28 × 2–8 µm, hyaline, thin to relatively thick-walled, with cell walls smooth; inner layer up to 15 µm thick, *textura porrecta* composed of hyaline hyphae up to 5 µm wide. Medullary excipulum up to 100 µm thick, composed of hyaline hyphae forming *textura intricata*; hyphae up to 3 µm wide. Hairs straight or irregularly curved, cylindrical, sometimes branched, up to 125 × 2.5–3.0 µm, hyaline, completely granulate, thin-walled; lacking crystals or resinous materials; apex usually equipped with hyaline apical amorphous materials. Asci (88–)92–101(–106) × 6–7.3(–8) µm (av. 96 ± 4.5 × 6.7 ± 0.6 µm, n = 18), 8-spored, thick-walled, cylindrical-clavate, arising from ascogenous hyphae branching several times; pore blue in MLZ without 3% KOH pretreatment; croziers absent at the basal septa. Ascospores (24–)26.7–34.5(–39) × (1.8–)1.9–2.3(–2.5) µm (av. 31 ± 3.9 × 2.1 ± 0.2 µm, n = 18), Q = (11–)12.5–17(–20)




**Figure 7.** *Erioscyphella insulae* TNS-F-39720 (Holotype) **A** dried apothecia **B** a pure culture on PDA (NBRC 114459) **C** asci **D** ascus pore MLZ (+) **E** ascospores **F** ascogenous hyphae **G** paraphyses **H** layer structures of excipulum **H1** medullary excipulum **H2** inner layer of ectal excipulum composed of hyphae **H3** outer layer of ectal excipulum composed of *textura angularis* **I, J** hairs with apical amorphous materials. Mounted in CB/LA (**C, E–J**), MLZ (**D**). Scale bars: 1 mm (**A**); 10  $\mu$ m (**A–J**).

(av.  $14.7 \pm 2.3$ ,  $n = 18$ ), showing various shapes and lengths, usually long fusiform and sometimes hypsiloid or sigmoid due to bending of both ends, sometimes swelling or constricted irregularly, aseptate or one- to three-septate (usually one-septate). Paraphyses straight, narrowly lanceolate, up to  $2.5 \mu$ m wide, septate, exceeding the asci up to  $7.5 \mu$ m.

**Culture characteristics.** Colony of NBRC 114445/TNS-F-26500 and NBRC 114459/TNS-F-39720 on PDA relatively thick-planar, pruinose, white to cream, ivory at the margin, pale sepia. Sectors and zonation absent. Aerial mycelium white to pale ocher, mainly developed except in the margin, not forming mycelial strands. Soluble pigment amber colored produced at the center. Margin unclear, flat and immersed into agar, radially undulate. Anamorph not seen.

**Distribution.** JAPAN (Bonin Islands, Yaeyama Islands).

**Notes.** This fungus resembles *Lachnum nothofagi* (Dennis) Spooner in the size and shape of apothecia, ascospores, asci, and hairs. However, *E. insulae* has completely hyaline hairs and ectal excipulum, and hairs are equipped with apical materials (Fig. 7J, 8A), whereas *L. nothofagi* has partly to totally brown hairs and ectal excipulum (Spooner 1987). *Lachnum nothofagi* is currently known only from New Zealand and Australia and mainly arises from *Nothofagus* spp., which are native in the southern hemisphere (Spooner 1987).




**Figure 8.** *Erioscyphella insulae* TNS-F-39720 (Holotype) **A** expansion of a vertical section of an apothecium **B** ascospores **C** apothecium **D** vertical section of an apothecium **E** asci **F** paraphyses **G** layer structures of excipulum.

***Erioscyphella otanii* Tochihara, sp. nov.**

MycoBank No: 835704

Figs 9, 10

**Diagnosis.** Characterized by pure white minute apothecia (< 0.3 mm in diameter) unlike *L. diminutum* with rather colored apothecia, and smaller asci compared to similar species *Lachnum minutum*.

**Holotype.** JAPAN, Hokkaido, Horonobe, Toikambetsu, Teshio Experimental Forest, Field Science Center for Northern Biosphere, Hokkaido University, 44.993978, 142.130125, ca 400 m, 11 Jul. 2018, on fallen leaves of *Sasa senanensis*, Y.Tochihara & K.Kaneko (TNS-F-81472).

**GenBank/UNITE no. ex holotype.** LC669471/UDB0779083 (ITS), LC533179 (LSU), LC533286 (mtSSU), LC533226 (RPB2).

**Other specimen examined.** JAPAN, Hokkaido, Sapporo, Mt. Moiwa, 43.024718, 141.318427, ca 530 m, 21 Jun. 1965, on fallen leaves of *Sasa kurilensis*, Y.Otani (TNS-F-50482, in poor condition).

**Etymology.** Referring to the name of Dr Yoshio Otani, the first discoverer of this species.

**Japanese name.** Kita-sasaba-hina-no-chawantake.




**Figure 9.** *Erioscyphella otanii* TNS-F-81472 (Holotype) **A** dried apothecia **B** pure culture on PDA (NBRC 114476) **C** asci **D** ascus pore MLZ (+) **E** ascospore **F** paraphyses **G** a hair **H** hair-apex with a apical amorphous material **I** ectal excipular cells. Mounted in CB/LA (**C**, **E–I**), MLZ (**D**). Scale bars: 0.5 mm (**A**); 10  $\mu$ m (**C–I**).

**Description.** Apothecia scattered, superficial, minute, 0.1–0.3 mm in diameter, at first spherical and later urceolate, having well-developed stipes, up to 0.3 mm high, pure white, externally covered with short white hairs, never colored brown. Disc concave, almost enclosed by an incurving margin when fresh and dry, cream to pale yellow when dry (not observed when fresh). Ectal excipulum *textura prismatica* like stone pavings arranged in rows, 3–25  $\times$  3–8  $\mu$ m, hyaline, relatively thick-walled; cell walls smooth. Medullary excipulum *textura intricata*; hyphae up to 2.5  $\mu$ m wide. Hairs straight, cylindrical or tapering toward the apices, up to 60  $\mu$ m long, up to 5  $\mu$ m wide near the bases and 2.5–3.0  $\mu$ m wide near the apices, arising from swollen ectal excipular cells, hyaline, up to 3-septate (usually 1- or 2-septate), thin-walled, completely granulated; granules dense near the apices and coarse toward the bases; apex sometimes with a hyaline and inconspicuous apical amorphous materials not dissolved with CB/LA, lacking any crystals or resinous materials. Asci (33–)34–38.8(–41)  $\times$  4–5  $\mu$ m (av.  $37 \pm 2.2 \times 4.4 \pm 0.4 \mu$ m,  $n = 15$ ), 8-spored, cylindrical-clavate, relatively thick-walled; pore blue in MLZ without 3% KOH pretreatment; croziers absent at the basal septa. Ascospores (11.5–)12.3–14.6(–15)  $\times$  (1.2–)1.36–1.7(–1.8)  $\mu$ m (av.  $13.4 \pm 1.2 \times 1.6 \pm 0.2 \mu$ m,  $n = 15$ ),  $Q = (6.7\text{--})7.8\text{--}9.6(10.8)$  (av.  $8.7 \pm 0.9$ ,  $n = 15$ ),




**Figure 10.** *Erioscyphella otanii* TNS-F-81472 (Holotype) **A** ascospores **B** apothecium **C** vertical section of an apothecium **D** hairs with cap-like structures arising from ectal excipular cells **E** expansion of a vertical section of an apothecium **F** paraphyses **G** asci.

fusiform, aseptate. Paraphyses straight, narrowly lanceolate to lanceolate, up to 2.5 µm wide, septate, exceeding the asci up to 10 µm.

**Culture characteristics.** Colony of NBRC 114476/TNS-F-81472 on PDA flat, partially protruding and forming mycelial mass, divided into two sectors. One sector flat, wooly to velvety, white to cream; dark ocher from the reverse. The other sector with wooly context, white and partly yellow; pale ocher from the reverse. Aerial mycelia developed throughout the colony, white, sparse to cottony, not forming mycelium strands. Margin distinct, flat and immersed into the agar. Soluble pigment absent. Asexual morph absent.

**Distribution.** JAPAN (Hokkaido; subarctic zone).

**Notes.** *Erioscyphella otanii* was first collected and documented by Otani (1967) under the misapplied name *Dasyscyphus diminutus* (TNS-F-50482). Based on the description, we concluded that the specimen was the same species as TNS-F-81472. The present species is very similar to *Lachnum diminutum* (Roberge ex Desm.) Rehm in the minute apothecia, ascospore size, and narrow paraphyses; however, *E. otanii* is pure white when fresh and dry (Fig. 9A, in dried state) and occurs on bamboo leaves, while *L. diminutum* is somewhat brown in the exterior parts of apothecia and occurs on sheaths of *Juncus* spp. (Dennis 1949). In the mature state, the apothecia of *E. otanii* become urceolate (Fig. 9A and Fig. 10B), whereas the apothecia of *L. diminutum* are flat (Dennis 1949). The ITS sequence of TNS-F-81472 showed low similarity (< 80%) with that of *L. diminutum* collected in France (GenBank accession number: MH857306). Based on the French sequence, *L. diminutum* is phylogenetically a good *Lachnum*.

The appearance of *E. otanii* is also similar to that of the graminicolous species *Lachnum minutum* W.Y. Zhuang and M. Ye documented in China (Ye and Zhuang 2003). *Erioscyphella otanii* is distinguished from *L. minutum* in having smaller asci, although DNA sequences of the species are not available.

***Erioscyphella papillaris* Tochihara, sp. nov.**

MycoBank No: 835705

Figs 11, 12

**Diagnosis.** Characterized by protruding papillary hairs with hyaline apical amorphous materials.

**Holotype.** JAPAN, Gunma, Minakami, Yubiso, Mt. Tanigawadake, 36.064014, 141.344653, ca 710 m, 16 Jul. 2017, on both sides of a fallen leaf of bamboo, Y.Tochihara (TNS-F-81272).




**Figure 11.** *Erioscyphella papillaris* TNS-F-81272 (Holotype) **A** dried apothecia **B** pure culture on PDA (NBRC 113937) **C** Ascus arising from ascogenous hyphae **D** an ascus **E** ascus pore iodine reactions **E1** MLZ (-) without 3% KOH pretreatment **E2** MLZ (-) with 3% KOH pretreatment **E3** IKI (+) without 3% KOH pretreatment **F** paraphysis **G** ascospores with guttules **H** ectal excipulum **I** hair-apex with a apical amorphous material **J** hairs. Mounted in CB/LA (**C, D, F-J**), MLZ (**E1, E2**), IKI (**E3**). Scale bars: 0.5 mm (**A**); 10  $\mu$ m (**C-J**).




**Figure 12.** *Erioscyphella papillaris* TNS-F-81272 (Holotype) **A** apothecium **B** vertical section of an apothecium **C** ascospores **D** expansion of an vertical section of an apothecium **E** ectal excipular cells **F** asci **G** paraphyses **H** hairs with cap-like structures.

**GenBank/UNITE no. ex holotype.** LC669473/UDB0779085 (ITS), LC533161 (LSU), LC533285 (mtSSU), LC533204 (RPB2).

**Etymology.** Referring to papillate hair apices.

**Japanese name.** Sasaba-hina-no-chawantake.

**Description.** Apothecia gregarious, superficial, minute, 0.1–0.3 mm in diameter, short-stipitate, up to 0.25 mm high, externally densely covered with pure white short hairs. Disc concave, white to lemon yellow when fresh and dry. Ectal excipulum *textura prismatica* composed of cuboid cells,  $3\text{--}13 \times 2.5\text{--}7\ \mu\text{m}$ , hyaline, thin-walled, lacking carotenoid pigments; cell walls smooth. Medullary excipulum *textura intricata* of hyaline hyphae up to  $3\ \mu\text{m}$  wide. Hairs straight, cylindrical,  $45\text{--}75 \times 3\text{--}5\ \mu\text{m}$ , 2–3-septate, hyaline, totally granulate, thin-walled, arising from swollen cells; apical cells rather longer than other cells,  $30\text{--}40\ \mu\text{m}$  long, with papillate at the apex, sometimes swelling,

equipped with hyaline and globose apical amorphous materials not dissolved with CB/LA, lacking any crystals or resinous matters. Asci (59–)59.8–66(–69)  $\times$  (7.5–)7.6–8.3(–9)  $\mu\text{m}$  (av.  $63 \pm 2.9 \times 8.0 \pm 0.4 \mu\text{m}$ ,  $n = 16$ ), 8-spored, cylindrical-clavate; pore inamyloid with MLZ without 3% KOH pretreatment, faint blue with MLZ with 3% KOH pretreatment, dark blue with IKI with and without KOH pretreatment; vesicle apparatus inverted-v-shaped present near the apices; croziers absent at the basal septa; base sympodially branched. Ascospores (16–)17.5–21.7(–24)  $\times$  (2–)2.3–2.8(–3)  $\mu\text{m}$  (av.  $20 \pm 2.1 \times 2.6 \pm 0.3 \mu\text{m}$ ,  $n = 20$ ),  $Q = (6.4\text{--})6.8\text{--}8.9(9.8)$  (av.  $7.8 \pm 1.0$ ,  $n = 20$ ), fusiform, aseptate, or one-septate (rarely two-septate), filled with hyaline oil drops. Paraphyses straight, cylindrical, up to 3  $\mu\text{m}$  wide, septate, containing small hyaline lipid bodies, equal or scarcely exceeding the asci.

**Culture characteristics.** Colony of NBRC 113937/TNS-F-81272 on PDA divided into two semicircular zones. The first zone umbonate, pruinose, white, producing white aerial mycelia densely, presenting wooly appearance; margin distinct, entire, flat. The second zone flat, glutinous, white to beige with concentric patterns, producing few aerial mycelia; margin entire, flat and immersed into agar, irregularly undulate. The reverse uniform unrelated to the zoning position, beige to pale dark brown throughout. Soluble pigment and asexual morph absent throughout the colony.

**Distribution.** JAPAN (Mt. Tanigawa). Currently known only from the type locality.

**Notes.** This species is similar to *Lachnum sclerotii* var. *microascum* in the dimension and shape of asci and ascospores, habitats, and inconspicuous ascus apex reaction in MLZ (Zhuang 2004). However, *E. papillaris* has ascospores containing conspicuous guttules in any mount (Fig. 11G) and filiform paraphyses rarely exceeding the asci (Fig. 11F, Fig. 12D, and Fig. 12G), whereas *L. sclerotii* var. *microascum* has non-guttulate asci and narrowly lanceolate to lanceolate paraphyses exceeding the asci by 15–18  $\mu\text{m}$  (Zhuang 2004). Although DNA sequences of *L. sclerotii* var. *microascum* are not available, we judged the present fungus as different from it, because the presence or absence of guttules in ascospores is a significant taxonomic character at the species level (Baral 2015).

Papillate hairs are also shown in the line drawings of *Lachnum gahniae* Spooner (Spooner 1987), suggesting the relationship of the present fungus to Australasian species. However, *L. gahniae* can be distinguished by having longer hairs, occurring on different substrates (leaves of Cyperaceae) and showing different ascal-iodine reactions (MLZ+) (Spooner 1987), although DNA sequences of *L. gahniae* are not available.

***Erioscyphella paralushanensis* Tochiara and Hosoya, sp. nov.**

MycoBank No: 839618

Figs 13, 14

**Diagnosis.** Characterized by throughout red apothecia occurring on bamboo sheaths. Similar to *E. lushanensis* in macro- and micromorphology and habitats, but has larger asci and ascospores.




**Figure 13.** *Erioscyphella paralushanensis* TNS-F-61920 (Holotype) **A** apothecia **B** pure culture on PDA (NBRC 114468) **C** ascus **D** ascal pore iodine reactions **D1** MLZ (faintly +) without 3% KOH pretreatment **D2** MLZ (+) with 3% KOH pretreatment **D3** IKI (+) without 3% KOH pretreatment **E** paraphysis **F** ascospores **G** ectal excipular cells **H** marginal section of an apothecium generating hairs **I** hairs with red resinous materials **J** apical amorphous materials of hairs. Mounted in CB/LA (**C**, **E–J**), MLZ (**D1**, **D2**), IKI (**D3**). Scale bars: 0.5 mm (**A**); 10  $\mu$ m (**C–J**).

**Holotype.** JAPAN, Shizuoka, Atami, Izusan, 35.128834, 139.051194, ca 620 m, 8 Jun. 2015, on fallen sheaths of *Pleioblastus argenteostriatus*, M.Nakajima (TNS-F-61920).

**GenBank/UNITE no. ex holotype.** LC669463/UDB0779075 (ITS), LC533141 (LSU), LC533267 (mtSSU), LC533220 (RPB2).

**Etymology.** Referring to the similarity with *E. lushanensis*.

**Japanese name.** Akage-hina-no-chawantake.

**Description.** Apothecia scattered, superficial, 0.7–1.5 mm in diameter, long-stipitate, up to 2.0 mm high, externally covered with dark-red hairs. Disc concave, cream to pale yellow. Ectal excipulum well-developed *textura prismatica* and partly *t. angularis*, 6–13  $\times$  2.0–2.5  $\mu$ m, hyaline, relatively thick-walled, with smooth walls. Medullary excipulum *textura intricata* of hyaline hyphae up to 2  $\mu$ m wide. Hairs straight, cylindrical, up to 160  $\mu$ m long, 2.0–3.0  $\mu$ m wide, pale brown but hyaline near the bases; hair cells narrowly septate, > 7  $\mu$ m long, covered by big and amber-colored granules; gran-




**Figure 14.** *Erioscyphella paralushanensis* TNS-F-61920 (Holotype) **A** apothecia **B** vertical section of an apothecium **C** expansion of a vertical section of an apothecium **D** asci **E** hairs **F** ectal excipulum **G** paraphyses **H** ascospores.

ules big and dense near the apices and smaller and sparse near the bases, up to 2  $\mu\text{m}$  in diameter near the apices, equipped with amber-colored resinous materials that dissolves in CB/LA at any position of hairs; apices with amber-colored apical amorphous materials, lacking any crystals. Asci (59–)61.4–70.2(–73)  $\times$  (4.5–)4.7–5.6(–6)  $\mu\text{m}$  (av.  $65.8 \pm 4.4 \times 5.2 \pm 0.4 \mu\text{m}$ ,  $n = 15$ ),  $Q = (11.5\text{--})12\text{--}13.6(\text{--}14.6)$  (av.  $12.8 \pm 0.8$ ,  $n = 15$ ), 8-spored, cylindrical-clavate; pore faintly blue in MLZ without 3% pretreatment, clear blue in MLZ with 3% KOH pretreatment and IKI without 3% KOH pretreatment. Ascospores (14–)15.8–20.7(–22)  $\times$  (1.5–)1.7–2.0  $\mu\text{m}$  (av.  $18.2 \pm 2.5 \times 1.8 \pm 0.2 \mu\text{m}$ ,  $n = 15$ ),  $Q = (7.5\text{--})8.7\text{--}11.2(\text{--}12.6)$  (av.  $9.9 \pm 1.3$ ,  $n = 15$ ), septate, sometimes bent to U-shaped or S-shaped, containing conspicuous guttules; guttules hyaline but sometimes red. Paraphyses straight, up to 2  $\mu\text{m}$  wide, septate, exceeding the asci 5–10  $\mu\text{m}$ , initially cylindrical to clavate, later becoming narrowly lanceolate.

**Culture characteristics.** Colony of NBRC 114468/TNS-F-61920 on PDA flat, sparse, dendritically spread. Context wooly, ocher to pale buff, dark buff from the reverse. Sectors and zonation absent. Aerial mycelium ocher to pale buff, dense cottony, developed near the center, forming white mycelium strands; margin distinct, flat and partly immersed into the agar. Asexual morph absent. Soluble pigments present, buff, dyeing agar without colony pale buff.

**Distribution.** JAPAN (Shizuoka). Currently known only from the type locality.

**Notes.** *Erioscyphella paralushanensis* is closely related to *E. lushanensis* in having red hairs (Fig. 13I) and the ectal excipulum composed of well-developed rectangular cells

in common (Fig. 13H, Fig. 14C, and Fig. 14F) (Zhuang and Wang 1998a). Compared with *E. lushanensis*, *E. paralushanensis* has slightly larger asci, ascospores and hairs. Red guttules in ascospores were observed only in *E. paralushanensis* (Fig. 13F). In this study, we proposed the present fungus as a new species, because species delimitation analyses based on ITS sequences strongly supported that *E. paralushanensis* is different from *E. lushanensis* (Fig. 3).

***Erioscyphella sasibrevispora* Tochihara & Hosoya, sp. nov.**

MycoBank No: 835706

Figs 15, 16

**Diagnosis.** Characterized by wooly appearance and yellow to orange discs, and distinguished from similar species *Lachnum novoguineense* var. *yunnanicum* in having shorter ascospores.

**Holotype.** JAPAN, Hokkaido, Tomakomai, Utonai, 42.705314, 141.7346, ca 10 m, 16 Jun. 2018, on fallen sheaths of *Sasa nipponica*, Y.Tochihara & T.Hosoya (TNS-F-81401).

**GenBank/UNITE no. ex holotype.** LC669470/UDB0779082 (ITS), LC533174 (LSU), LC533269 (mtSSU), LC533217 (RPB2).

**Other specimen examined.** JAPAN, Gunma, Higashiagatsuma, 36.562253, 138.724139, ca 1330 m, 6 Jun. 2017, on fallen sheaths of *Sasa veitchii*, Y.Tochihara & T.Hosoya (TNS-F-80399, in bad condition).

**Etymology.** “sasi” means bamboo [host plants] and “brevispora” means shorter ascospores compared to *L. novoguineense* var. *yunnanicum*.

**Japanese name.** Sasa-no-youmou-chawantake.

**Description.** Apothecia gregarious, superficial, 0.6–1.3 mm in diameter, short-stipitate, up to 0.8 mm high, pure white, externally covered with long white hairs. Disc concave, yellow to pale orange when fresh and dry. Ectal excipulum *textura prismatica* to *t. angularis*, 3–16 × 2–10 µm, hyaline, thin-walled; surface smooth. Medullary excipulum *textura intricata* of hyaline hyphae up to 2 µm wide. Hairs straight, delicate, cylindrical with relatively acute apices, up to 190 × 2–3 µm, hyaline, totally granulate, thin-walled; apical cell a little longer than other cells, lacking any crystals, resinous materials, or apical amorphous materials. Asci (79–)82.5–90(–95) × (6–)6.6–8.1(–9) µm (av. 86 ± 4.0 × 7.4 ± 0.8 µm, n = 15), 8-spored, cylindrical-clavate; lateral parts sometimes swelling irregularly; pore blue in MLZ without 3% KOH pretreatment; croziers with perforation present at the basal septa. Ascospores (26–)27.9–36.1(–39) × (1.5–)1.7–2 µm (av. 32 ± 4.1 × 1.8 ± 0.2 µm, n = 17), Q = (13–)15–19.7(–21) (av. 17.5 ± 2.3, n = 17), long fusiform, usually 3-septate, rarely 0- to 2-septate (only observed in TNS-F-81401 because TNS-F-80399 was immature). Paraphyses straight, lanceolate, 2.5–4 µm wide, densely septate, exceeding the asci up to 15 µm. Note that the description is solely based on the holotype because another examined specimen TNS-F-80399 was in bad condition.




**Figure 15.** *Erioscyphella sasibrevispora* TNS-F-81401 (Holotype, **A–F, H–J**). *Lachnum novoguineense* var. *yunnanicum* TNS-F-16442 (**G**) **A** dried apothecia **B** a pure culture on PDA (NBRC 114475) **C** ectal excipular cells **D** ascus **E** an ascus pore MLZ (+) **F** ascus base with a perforated crozier **G** ascus base with a perforated crozier **H** septated paraphyses **I** ascospores **J** vertical section through the apothecium. Mounted in CB/LA (**D, F–J**), MLZ (**E**). Scale bars: 1 mm (**A**); 10  $\mu$ m (**C–J**).

**Culture characteristics.** Colony of NBRC 114475/TNS-F-81401 on PDA wrinkled. Context cottony and partially funiculose, white, turning ocher at the center; almost ocher except for the white margin from the reverse. Sectors and zonation absent. Aerial mycelium developed throughout the colony, concolous, forming mycelium strands. Margin indistinct, flat and immersed into agar. Soluble pigment absent. Asexual morph absent.

**Distribution.** JAPAN (cool-temperate zone, subarctic zone).

**Notes.** *Erioscyphella sasibrevispora* is closely related to *L. novoguineensis* var. *yunnanicum* (TNS-F-16442, 16642) (Fig. 1) and occurs in the same habitats (that is, bamboo sheaths) but has shorter asci and ascospores. The ascus bases of the two species are very characteristic, in that they have croziers with perforations (Fig. 15G and Fig. 16E). In Lachnaceae, this type of crozier has only been reported in *Lachnel-*




**Figure 16.** *Erioscyphella sasibrevispora* TNS-F-81401 (Holotype **A–D, F, G**). *Lachnum novoguineense* var. *yunnanicum* TNS-F-16642 (**E**) **A** apothecium **B** vertical section of an apothecium **C** ascospores **D** asci (with basal structures sometimes with perforation) **E** ascus base arising from a crozier with perforation **F** paraphyses **G** ectal excipular cells **H** hairs.

*lula* (Baral 1984). Additionally, both species exceptionally lack any hair materials in *Erioscyphella*.

The tropical species *E. bambusina* and *Lachnum albidum* var. *americanum* (Dennis) W.Y. Zhuang also occur on bamboo sheaths. However, compared with the present fungus, the former has smaller ascospores and filiform paraphyses (Dennis 1954), and the latter has extremely large asci and ascospores (Dennis 1960). In cool-temperate to subarctic zones, *L. asiaticum* and *Lachnum sasae* Raitv. occur on bamboo sheaths (Otani 1967; Raitviir 1985), but their ascospores are much shorter than those of the present fungus.

The woolly appearance and yellow disc of this species (Fig. 15A) resemble those of *Capitotricha rubi* (Bres.) Baral; however, microscopic observations easily distinguish the two species.

***Erioscyphella sinensis* (Z.H. Yu and W.Y. Zhuang) Sasagawa, Tochihara & Hosoya, comb. et stat. nov.**

MycoBank No: 835709

≡ *Lachnum mapirianum* var. *sinense* Z.H. Yu and W.Y. Zhuang, Nova Hedwigia 74(3-4): 422 (2002).

**Diagnosis.** Occurring on fallen leaves of *Quercus* spp. or *Castanopsis* spp. in early summer and having needle-like ascospores.

**Japanese name.** Shii-Kashi-hina-no-chawantake-modoki.

**Specimen examined.** JAPAN, Ibaraki, Tsukuba, Mt. Tsukuba, 36.228539, 140.103504, ca 870 m, 23 Jun. 2007, on fallen leaves of *Castanopsis sieboldii*, R.Sasagawa (TNS-F-16841). JAPAN, Ibaraki, Tsukuba, Amakubo, Tsukuba Botanical Garden, 36.101472, 140.110944, ca 20 m, 15 Jun. 2007, on fallen leaves of *C. sieboldii*, R.Sasagawa (TNS-F-16838). JAPAN, Tottori, Yonago, Yonago Castle, 35.42437, 133.325472, ca 50 m, 3 Jun. 2018, on fallen leaves of *C. sieboldii*, Y.Tochihara (TNS-F-81383).

**Distribution.** CHINA (Hainan, Yunnan; Yu and Zhuang 2003). JAPAN (warm-temperate zone).

**Notes.** The present fungus was treated as *Lachnum* sp. 13 by Hosoya et al. (2010). This fungus occurs in the same habitats as *E. hainanensis*, but it is easily distinguished in having longer and needle-like ascospores. *Erioscyphella sinensis* resembles *L. mapirianum* in the shape of ascospores, but the two species are different in that *L. mapirianum* has long slender apothecial stipes, larger asci, longer ascospores, and wider paraphyses.

In the present study, we transferred this fungus to *Erioscyphella* and upgraded it from variety to species level, because this fungus is not phylogenetically related to '*L. mapirianum*' (Fig. 1). The presence of apical amorphous materials of hairs was confirmed in this study (Suppl. material 1: Fig. S2).

## Acknowledgements

We thank Dr Shimpei Hiruta at the National Museum of Nature and Science for his kind support in the species delimitation analyses. We also thank Dr Toshimitsu Fukiharu at the Natural History Museum and Institute, Chiba, Ms Michiru Fujisaki and Rei Sasagawa at the Faculty of Life and Environmental Sciences, University of Tsukuba, and Mr Minoru Nakajima at Kanagawa Kinoko no Kai for collecting and donating their significant fungal specimens to TNS.

## References

- Abarenkov K, Tedersoo L, Nilsson RH, Vellak K, Saar I, Veldre V, Parmasto E, Proulx M, Aan A, Ots M, Kurina O, Ostonen I, Jõgeva J, Halapuu S, Põldmaa K, Toots M, Truu J, Larsson K-H, Kõljalg U (2010) PlutoF – a web based workbench for ecological and taxonomic

- research, with an online implementation for fungal ITS sequences. *Evolutionary Bioinformatics* 6: 189–196. <https://doi.org/10.4137/EBO.S6271>
- Akaike H (1974) A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19(6): 716–723. <https://doi.org/10.1109/TAC.1974.1100705>
- Baral HO (1984) Taxonomische und ökologische Studien über die Koniferen bewohnenden europäischen Arten der Gattung *Lachnellula* Karsten. *Beiträge zur Kenntnis der Pilze Mitteleuropas* 1: 143–156. [in German]
- Baral HO (2009) Iodine reaction in Ascomycetes: why is Lugol's solution superior to Melzer's reagent? <http://www.gbif-mycology.de/HostedSites/Baral/IodineReaction.htm> [Accessed on: 2021-03-15]
- Baral HO (2015) *Hymenoscyphus menthae*, *H. macroguttatus* and *H. scutula*, a comparative taxonomic study emphasizing the value of spore guttulation and croziers. *Ascomycete.org* 7(6): 255–287. <https://doi.org/10.25664/art-0147>
- Baral HO, Krieglsteiner GJ (1985) Bausteine zu einer Askomyzeten-Flora der Bundesrepublik Deutschland. In: *Süddeutschland gefundene inoperculte Diskomyceten mit taxonomischen, ökologischen, chorologischen Hinweisen und einer Farbtafel*. Beiheften zur Zeitschrift für Mykologie 6: 1–160. [in German]
- Begerow D, Nilsson H, Unterseher M, Maier W (2010) Current state and perspectives of fungal DNA barcoding and rapid identification procedures. *Applied Microbiology and Biotechnology* 87: 99–108. <https://doi.org/10.1007/s00253-010-2585-4>
- Biczok R, Bozsoky P, Eisenmann P, Ernst J, Ribizel T, Scholz F, Trefzer A, Weber F, Hamann M, Stamatakis A (2018) Two C++ libraries for counting trees on a phylogenetic terrace. *Bioinformatics* 34(19): 3399–3401. <https://doi.org/10.1093/bioinformatics/bty384>
- Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R, Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, Koester I, Kosciolk T, Kreps J, Langille MGI, Lee J, Ley R, Liu YX, Loftfield E, Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik AV, Metcalf JL, Morgan SC, Morton JT, Naimey AT, Navas-Molina JA, Nothias LF, Orchanian SB, Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB, Rivers A, Robeson MS 2<sup>nd</sup>, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD, Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft JJJ, Vargas F, Vázquez-Baeza Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang M, Warren J, Weber KC, Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG (2019) Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology* 37: 852–857. <https://doi.org/10.1038/s41587-019-0209-9>
- Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, Heled J, Jones G, Kühnert D, Maio ND, Matschiner M, Mendes FK, Müller NE, Ogilvie HA, Plessis L, Poppinga A, Rambaut A, Rasmussen D, Siveroni I, Suchard MA, Wu C-H,

- Xie D, Zhang C, Stadler T, Drummond AJ (2019) BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLOS Computational Biology* 15(4): e1006650. <https://doi.org/10.1371/journal.pcbi.1006650>
- Cantrell SA, Hanlin RT (1997) Phylogenetic relationships in the family Hyaloscyphaceae inferred from sequences of ITS regions, 5.8S ribosomal DNA and morphological characters. *Mycologia* 89(5): 745–755. <https://doi.org/10.1080/00275514.1997.12026841>
- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25(15): 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>
- Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T (2019) ModelTest-NG: A new and scalable tool for the selection of DNA and protein evolutionary models. *Molecular Biology and Evolution* 37(1): 291–294. <https://doi.org/10.1093/molbev/msz189>
- Dennis RWG (1949) A revision of the British Hyaloscyphaceae with notes on related European species. *Mycological Papers* 32: 1–97.
- Dennis RWG (1954) Some inoperculate discomycetes of tropical America. *Kew Bulletin* 9(2): 289–348. <https://doi.org/10.2307/4114399>
- Dennis RWG (1960) Fungi venezuelani: III. *Kew Bulletin* 14(3): 418–458. <https://doi.org/10.2307/4114758>
- Dharne CG (1965) Taxonomic investigations on the discomycetous genus *Lachnellula* Karst. *Phytopathologische Zeitschrift* 53(2): 101–144. <https://doi.org/10.1111/j.1439-0434.1965.tb02194.x>
- Ekanayaka AH, Hyde KD, Gentekaki E, McKenzie EHC, Zhao Q, Bulgakov TS, Camporesi E (2019) Preliminary classification of Leotiomycetes. *Mycosphere* 10: 310–489. <https://doi.org/10.5943/mycosphere/10/1/7>
- Felsenstein J (1984) Distance methods for inferring phylogenies: a justification. *Evolution* 38: 16–24. <https://doi.org/10.1111/j.1558-5646.1984.tb00255.x>
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791. <https://doi.org/10.2307/2408678>
- Fujisawa T, Barraclough TG (2013) Delimiting species using single-locus data and the Generalized Mixed Yule Coalescent approach: a revised method and evaluation on simulated data sets. *Systematic Biology* 62(5): 707–724. <https://doi.org/10.1093/sysbio/syt033>
- Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes-application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2(2): 113–118. <https://doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- GBIF (2018) Adding sequence-based identifiers to backbone taxonomy reveals ‘dark taxa’ fungi. <https://www.gbif.org/news/2LrgV5t3ZuGeU2WIymSEuk/adding-sequence-based-identifiers-to-backbonetaxonomy-reveals-dark-taxa-fungi> [Accessed on: 2020-02-10]
- Gelardi M, Vizzini A, Ercole E, Horak E, Ming Z, Li TH (2015) Circumscription and taxonomic arrangement of *Nigroboletus roseonigrescens* Gen. Et Sp. Nov., a new member of Boletaceae from tropical South-Eastern China. *PloS ONE* 10: e0134295. <https://doi.org/10.1371/journal.pone.0134295>
- Guatimosim E, Schwartsburd PB, Barreto RW, Crous PW (2016) Novel fungi from an ancient niche: cercosporoid and related sexual morphs on ferns. *Persoonia* 37: 106–141. <https://doi.org/10.3767/003158516X690934>

- Haines JH (1980) Studies in the Hyaloscyphaceae. I: Some species of *Dasyscyphus* on tropical ferns. *Mycotaxon* 11(1): 189–216.
- Haines JH (1992) Studies in the Hyaloscyphaceae. VI: The genus *Lachnum* (ascomycetes) of the Guayana Highlands. *Nova Hedwigia* 54: 97–112.
- Haines JH, Dumont KP (1983) Studies in the Hyaloscyphaceae II: *Proliferodiscus*, A new genus of Arachnopezizoideae. *Mycologia* 75(3): 535–543. <https://doi.org/10.1080/00275514.1983.12023717>
- Haines JH, Dumont KP (1984) Studies in the Hyaloscyphaceae III: The long-spored, lignicolous species of *Lachnum*. *Mycotaxon* 19: 1–39.
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Han JG, Hosoya T, Sung GH, Shin HD (2014) Phylogenetic reassessment of Hyaloscyphaceae sensu lato (Helotiales, Leotiomycetes) based on multigene analyses. *Fungal Biology* 118: 150–167. <https://doi.org/10.1016/j.funbio.2013.11.004>
- Hansen K, LoBuglio KF, Pfister DH (2005) Evolutionary relationships of the cup-fungus genus *Peziza* and Pezizaceae inferred from multiple nuclear genes: RPB2,  $\beta$ -tubulin, and LSU rDNA. *Molecular Phylogenetics and Evolution* 36(1): 1–23. <https://doi.org/10.1016/j.ympev.2005.03.010>
- Hosaka K, Castellano MA (2008) Molecular phylogenetics of Geastrales with special emphasis on the position of *Sclerogaster*. *Bulletin of the National Science Museum. Series B, Botany* 34(4): 161–173. [https://www.kahaku.go.jp/research/publication/botany/download/34\\_4/BNMNS\\_B340403.pdf](https://www.kahaku.go.jp/research/publication/botany/download/34_4/BNMNS_B340403.pdf) [Accessed on: 2021-03-21]
- Hosoya T (2021) Systematics, ecology, and application of Helotiales: Recent progress and future perspectives for research with special emphasis on activities within Japan. *Mycoscience* 62(1): 1–9. <https://doi.org/10.47371/mycosci.2020.05.002>
- Hosoya T, Saito Y, Sasagawa R (2013) Enumeration of remarkable Japanese discomycetes (7): Notes on one operculate discomycete and one inoperculate discomycete. *Bulletin of the National Science Museum. Series B, Botany* 39(4): 151–158. [https://www.kahaku.go.jp/research/publication/botany/download/39\\_4/BNMNS\\_B39-4\\_151-158.pdf](https://www.kahaku.go.jp/research/publication/botany/download/39_4/BNMNS_B39-4_151-158.pdf) [Accessed on: 2020-03-12]
- Hosoya T, Sasagawa R, Hosaka K, Gi-Ho S, Hirayama Y, Yamaguchi K, Toyama K, Kakishima M (2010) Molecular phylogenetic studies of *Lachnum* and its allies based on the Japanese material. *Mycoscience* 51(3): 170–180. <https://doi.org/10.1007/S10267-009-0023-1>
- Index Fungorum (2021) Index Fungorum. <http://www.indexfungorum.org/names/Names.asp> [Accessed on: 2021-04-15]
- Johnston PR, Quijada L, Smith CA, Baral HO, Hosoya T, Baschien C, Pärtel K, Zhuang WY, Haelewaters D, Park D, Carl S, López-Giráldez F, Wang Z, Townsend JP (2019) A multigene phylogeny toward a new phylogenetic classification of Leotiomycetes. *IMA Fungus* 10: e1. <https://doi.org/10.1186/s43008-019-0002-x>
- Kanouse BB (1941) New and unusual species of discomycetes. *Mycologia* 33: 461–467. <https://doi.org/10.1080/00275514.1941.12020841>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>

- Kirschstein W (1938) Über neue, seltene und kritische Ascomyceten und Fungi imperfecti. I. *Annales Mycologici* 36(5–6): 367–400.
- Kõljalg U, Tedersoo L, Nilsson RH, Abarenkov K (2016) Digital identifiers for fungal species. *Science* 352(6290): 1182–1183. <https://doi.org/10.1126/science.aaf7115>
- Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AF, Bahram M, Bates ST, Bruns TD, Bengtsson-Palme J, Callaghan TM, Douglas B, Drenkhan T, Eberhardt U, Dueñas M, Grebenc T, Griffith GW, Hartmann M, Kirk PM, Kohout P, Larsson E, Lindahl BD, Lücking R, Martín MP, Matheny PB, Nguyen NH, Niskanen T, Oja J, Peay KG, Peintner U, Peterson M, Põldmaa K, Saag L, Saar I, Schüßler A, Scott JA, Senés C, Smith ME, Suija A, Taylor DL, Telleria MT, Weiss M, Larsson KH (2013) Towards a unified paradigm for sequence-based identification of fungi. *Molecular Ecology* 22: 5271–5277. <https://doi.org/10.1111/mec.12481>
- Kõljalg U, Nilsson HR, Schigel D, Tedersoo L, Larsson K-H, May TW, Taylor AFS, Jeppesen TS, Frøslev TG, Lindahl BD, Põldmaa K, Saar I, Suija A, Savchenko A, Yatsiuk I, Adojaan K, Ivanov F, Piirmann T, Pöhönen R, Zirk A, Abarenkov K (2020) The Taxon Hypothesis Paradigm – On the Unambiguous Detection and Communication of Taxa Microorganisms 8(12): e1910. <https://doi:10.3390/microorganisms8121910>
- Korf RP (1978) Nomenclatural and taxonomic notes on *Lasiobelonium*, *Erioscypha* and *Erioscyphella*. *Mycotaxon* 7(2): 399–406.
- Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A (2019) RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* 35(21): 4453–4455. <https://doi.org/10.1093/bioinformatics/btz305>
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution* 35(6): 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Lemoine F, Entfellner JBD, Wilkinson E, Correia D, Felipe MD, Oliveira TD, Gascuel O (2018) Renewing Felsenstein's phylogenetic bootstrap in the era of big data. *Nature* 556: 452–456. <https://doi.org/10.1038/s41586-018-0043-0>
- Liu YL, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16: 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Matheny PB (2005) Improving phylogenetic inference of mushrooms with RPB1 and RPB2 nucleotide sequences (Inocybe, Agaricales). *Molecular Phylogenetics and Evolution* 35: 1–20. <https://doi.org/10.1016/j.ympev.2004.11.014>
- Matheny PB, Wang Z, Binder M, Curtis JM, Lim YW, Nilsson RH, Hughes KW, Hofstetter V, Ammirati JF, Schoch CL, Langer E, Langer G, McLaughlin DJ, Wilson AW, Frøslev T, Ge ZW, Kerrigan RW, Slot JC, Yang ZL, Baroni TJ, Fischer M, Hosaka K, Matsuura K, Seidl MT, Vauras J, Hibbett DS (2007) Contributions of rpb2 and tef1 to the phylogeny of mushrooms and allies (Basidiomycota, Fungi). *Molecular Phylogenetics and Evolution* 43(2): 430–451. <https://doi.org/10.1016/j.ympev.2006.08.024>
- Miyoshi T, Ono Y, Shimizu S (2007) Occurrence of concave stem canker of citrus in Ehime prefecture [Japan] and detection of the pathogenic fungus *Lachnum abnorme* by PCR. *Japanese Journal of Phytopathology* 73(1): 9–14. <https://doi.org/10.3186/jjphytopath.73.9> [in Japanese]

- Nilsson RH, Abarenkov K, Veldre V, Nylinder S, De Wit P, Brosché S, Alfredsson JF, Ryberg M, Kristiansson E (2010). An open source chimera checker for the fungal ITS region. *Molecular Ecology Resources* 10(6): 1076–1081. <https://doi.org/10.1111/j.1755-0998.2010.02850.x>
- Nilsson RH, Tedersoo L, Ryberg M, Kristiansson E, Hartmann M, Unterseher M, Porter TM, Bengtsson-Palme J, Walker DM, de Sousa F, Gamper HA, Larsson E, Larsson KH, Kõljalg U, Edgar RC, Abarenkov K (2015) A comprehensive, automatically updated fungal ITS sequence dataset for reference-based chimera control in environmental sequencing efforts. *Microbes and Environments* 30(2): 145–150. <https://doi.org/10.1264/jsme2.ME14121>
- Otani Y (1967) Notes on some cup fungi of the Hyaloscyphaceae collected in Hokkaido, Japan. *Transactions of the Mycological Society of Japan* 8: 33–42.
- Perić B, Baral HO (2014) *Erioscyphella curvispora* spec. nov. from Montenegro. *Mycologia Montenegrina* 17: 89–104. <https://www.etis.ee/File/DownloadPublic/ab9f7b44-b094-4148-8508-a48479be0ba9?> [Accessed on: 2020-02-15]
- Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP, Hazell S, Kamoun S, Sumlin WD, Vogler AP (2006) Sequence-based species delimitation for the DNA taxonomy of undescribed insects. *Systematic Biology* 55(4): 595–609. <https://doi.org/10.1080/10635150600852011>
- Puillandre N, Brouillet S, Achaz G (2021) ASAP: assemble species by automatic partitioning. *Molecular Ecology Resources* 21(2): 609–620. <https://doi.org/10.1111/1755-0998.13281>
- Raitviir A (1985) Species Hyaloscyphacearum in *Sasa* spp. inventae. *Novosti Sistematiki Nizshikh Rastenii* 22: 157–162.
- Raitviir A (2002) A revision of the genus *Dasyscyphella* (Hyaloscyphaceae, Helotiales). *Polish Botanical Journal* 47(2): 227–241.
- Rambaut A (2018a) Tracer. Molecular evolution, phylogenetics and epidemiology, Edinburgh. <http://beast.community/tracer> [Accessed on: 2021-03-15]
- Rambaut A (2018b) FigTree. Molecular evolution, phylogenetics and epidemiology, Edinburgh. <http://tree.bio.ed.ac.uk/software/figtree/> [Accessed on: 2020-03-15]
- Rognes T, Flouri T, Nichols B, Quince C, Mahé F (2016) VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4: e2584. <https://doi.org/10.7717/peerj.2584>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Sanderson MJ, McMahon MM, Steel M (2011) Terraces in phylogenetic tree space. *Science* 333(6041): 448–450. <https://doi.org/10.1126/science.1206357>
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W, Fungal Barcoding Consortium (2012) Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *PNAS* 109(16): 6241–6246. <https://doi.org/10.1073/pnas.1117018109>
- Suková M (2005) A revision of selected material of lignicolous species of *Brunnipila*, *Capitotricha*, *Dasyscyphella* and *Neodasyscypha* from the Czech Republic. *Czech Mycology* 57(1–2): 139–172. <https://doi.org/10.33585/cmy.57108>

- Spooner BM (1987) Helotiales of Australasia: Geoglossaceae, Orbiliaceae, Sclerotiniaceae, Hyaloscyphaceae. *Bibliotheca Micologica* 116: 1–711.
- Swofford DL (2002) PAUP\*. Phylogenetic analysis using parsimony (\* and other methods). Version 4.0 b10. Sinauer Associates, Sunderland, MA.
- Tello S, Baral HO (2016) *Erioscyphella lunata* (Lachnaceae), a rare discomycete collected in Spain. *Ascomycete.org* 8(4): 157–162. <https://doi.org/10.25664/ART-0183>
- Tochihara Y, Hosoya T (2019) Three new species of *Incrucipulum* (Lachnaceae, Helotiales, Ascomycota) from Japan. *Phytotaxa* 403: 25–38. <https://doi.org/10.11646/phytotaxa.403.1.2>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- White T, Bruns TD, Lee A, Taylor JW (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Snisky JJ, White TJ (Eds) *PCR Protocols: A Guide to Methods and Applications*. Academic Press, San Diego, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Ye M, Zhuang WY (2003) New taxa of *Lachnum* (Helotiales, Hyaloscyphaceae) from temperate China. *Nova Hedwigia* 76(3–4): 443–450. <https://doi.org/10.1127/0029-5035/2003/0076-0443>
- Zhao M, Yuan LY, Guo DL, Ye Y, Da-Wa ZM, Wang XL, Ma FW, Chen L, Gu YC, Ding LS, Zhou Y (2018) Bioactive halogenated dihydroisocoumarins produced by the endophytic fungus *Lachnum palmae* isolated from *Przewalskia tangutica*. *Phytochemistry* 148: 97–103. <https://doi.org/10.1016/j.phytochem.2018.01.018>
- Zhao P, Zhuang WY (2011) Evaluation of ITS region as a possible DNA barcode for the genus *Lachnum* (Helotiales). *Mycosystema* 30(6): 932–937.
- Zhao YJ, Hosoya T, Baral HO, Hosaka K, Kakishima M (2012) *Hymenoscyphus pseudoalbidus*, the correct name for *Lambertella albida* reported from Japan. *Mycotaxon* 122: 25–41. <https://doi.org/10.5248/122.25>
- Zhuang WY (2004) New taxa of *Lachnum* (Ascomycetes, Helotiales) on bamboo and a key to the bambusicolous species of the genus. *Nova Hedwigia* 78(3–4): 425–433. <https://doi.org/10.1127/0029-5035/2004/0078-0425>
- Zhuang WY, Wang Z (1998a) Some new species and new records of discomycetes in China. VIII. *Mycotaxon* 66: 429–438.
- Zhuang WY, Wang Z (1998b) Discomycetes of tropical China. I. Collections from Hainan Island. *Mycotaxon* 67: 21–31.
- Zhang J, Kapli P, Pavlidis P, Stamatakis A (2013) A general species delimitation method with applications to phylogenetic placements. *Bioinformatics* 15(29): 2869–2876. <https://doi.org/10.1093/bioinformatics/btt499>
- Zoller S, Scheidegger C, Sperisen C (1999) PCR primers for the amplification of mitochondrial small subunit ribosomal DNA of lichen-forming ascomycetes. *The Lichenologist* 31: 511–516. <https://doi.org/10.1006/lich.1999.0220>

## Supplementary material 1

### Figure S1. ML trees

Authors: Yukito Tochihara, Tsuyoshi Hosoya

Data type: Image.

Explanation note: ML trees based on ITS (**A**), LSU (**B**), mtSSU (**C**) and RPB2 (**D**) constructed using MEGA X. Bootstrap values > 50% are indicated on branches and branches with MLBS > 70% are shown bold.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.87.73082.suppl1>

## Supplementary material 2

### Figure S2. Hair apices

Authors: Yukito Tochihara, Tsuyoshi Hosoya

Data type: Image.

Explanation note: Hair apices of members of Clade **A** *Erioscyphella abnormis* TNS-F-32163 **B** *E. abnormis* TNS-F-61773 **C** *E. brasiliensis* TNS-F-46419 **D** *E. sclerotii* TNS-F-26492 **E** ‘*Lachnum*’ *mapirianum* TNS-F-17245 **F** ‘*Lachnum*’ *palmae* TNS-F-17567 **F1** Hair with resinous matters **F2** Hair with apical amorphous material **G** ‘*Lachnum*’ *palmae* TNS-F-24600 **G1** Hair with a resinous matter **G2** Hair with apical amorphous materials **H** *E. hainanensis* TNS-F-80371 **I** *E. sinensis* TNS-F-80354. Mounted in CB/LA. Scale bars: 10 mm. Arrowheads show hair apical materials.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.87.73082.suppl2>

### Supplementary material 3

#### Figure S3. Result of the ASAP species delimitation analysis

Authors: Yukito Tochihara, Tsuyoshi Hosoya

Data type: Image.

Explanation note: The graph shows the distribution of ASAP scores according to partitioning results, and the phylogenetic tree shows the way of partitioning.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.87.73082.suppl3>

### Supplementary material 4

#### Figure S4. Result of the GMYC species delimitation analysis

Authors: Yukito Tochihara, Tsuyoshi Hosoya

Data type: Image.

Explanation note: Number with each node shows the support value that each cluster is an independent species.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.87.73082.suppl4>

### Supplementary material 5

#### Figure S5. ML best-scored phylogenetic tree based on concatenated dataset of ITS1, 5.8S, and ITS2 constructed by RAxML-NG

Authors: Yukito Tochihara, Tsuyoshi Hosoya

Data type: Image.

Explanation note: GenBank/UNITE accession number and TNS specimen number (if any) is shown for each taxon. MLBP > 50% were attached on branches.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.87.73082.suppl5>

## Supplementary material 6

### Figure S6. Results of PTP species delimitation analyses

Authors: Yukito Tochihara, Tsuyoshi Hosoya

Data type: Image.

Explanation note: Number with each node shows the probability of the likelihood that each cluster is an independent species. Clusters showed by red branches are regarded as species.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.87.73082.suppl6>